arXiv:1001.0755v1 [math.FA] 5 Jan 2010

Sharp estimates for the commutators of the Hilbert, Riesz transforms and the Beurling-Ahlfors operator on weighted Lebesgue spaces

Daewon Chung

January 5, 2010

Abstract

We prove that the operator norm on weighted Lebesgue space $L^2(w)$ of the commutators of the Hilbert, Riesz and Beurling transforms with a BMO function b depends quadratically on the A_2-characteristic of the weight, as opposed to the linear dependence known to hold for the operators themselves. It is known that the operator norms of these commutators can be controlled by the norm of the commutator with appropriate Haar shift operators, and we prove the estimate for these commutators. For the shift operator corresponding to the Hilbert transform we use Bellman function methods, however there is now a general theorem for a class of Haar shift operators that can be used instead to deduce similar results. We invoke this general theorem to obtain the corresponding result for the Riesz transforms and the Beurling-Ahlfors operator. We can then extrapolate to $L^p(w)$, and the results are sharp for $1 < p < \infty$. [1][2]

1 Introduction

We say w is a weight if it is positive almost everywhere and locally integrable. The norm of $f \in L^p(w)$ is

$$\|f\|_{L^p(w)} := \left(\int_{\mathbb{R}} |f(x)|^p w(x) dx \right)^{1/p} .$$

Helson and Szegö first found necessary and sufficient conditions for the boundedness of the Hilbert transform

$$Hf(x) = p.v. \frac{1}{\pi} \int \frac{f(y)}{x-y} dy$$

in weighted Lebesgue spaces in [11]. Hunt, Muckenhoupt, and Wheeden showed in [13] that a new necessary and sufficient condition for boundedness of the Hilbert transform in $L^p(w)$ is that the weight satisfies the A_p condition, namely:

$$[w]_{A_p} := \sup_I \langle w \rangle_I \langle w^{-1/(p-1)} \rangle_I^{p-1} < \infty \,, \tag{1.1}$$

where we denote the average over the interval I by $\langle \cdot \rangle_I$, and we take the supremum over all intervals in $\mathbb{R}$. After one year, Coifman and Fefferman extended in [5] the result to a larger class of convolution singular integrals with standard kernels. Recently, many authors have been interested in finding the

---

[1] Key words and phrases: Operator-weighted inequalities, Commutator, Hilbert transform, Paraproduct

[2] 2000 Mathematics Subject Classification. Primary 42A50; Secondary 47B47.

sharp bounds for the operator norms in terms of the A_p-characteristic $[w]_{A_p}$ of the weight. That is, one looks for a function $\phi(x)$, sharp in terms of its growth, such that

$$\|Tf\|_{L^p(w)} \leq C\phi\big([w]_{A_p}\big)\|f\|_{L^p(w)}\,.$$

For $T = M$, the Hardy-Littlewood maximal function, S. Buckley [3] showed that $\phi(x) = x^{1/(p-1)}$ is the sharp rate of growth for all $1 < p < \infty$. He also showed in [3] that $\phi(x) = x^2$ works for the Hilbert transform in $L^2(w)$. S. Petermichl and S. Pott improved the result to $\phi(x) = x^{3/2}$, for the Hilbert transform in $L^2(w)$ in [28]. More recently, S. Petermichl proved in [26] the linear dependence, $\phi(x) = x$, for the Hilbert transform in $L^2(w)$

$$\|Hf\|_{L^2(w)} \leq C[w]_{A_2}\|f\|_{L^2(w)}\,,$$

by estimating the operator norm of the dyadic shift. Linear bounds in $L^2(w)$ were also obtained by O. Beznosova for the dyadic paraproduct [2]. Most recently there are new proofs for the linear estimates in $L^2(w)$ for some operators including the Hilbert transform and the dyadic paraproduct, [17] and [7]. The conjecture is that all the Calderón-Zygmund singular integral operator obey linear bounds in weighted L^2. So far this is known only for the Beurling-Ahlfors operator [9], [30], the Hilbert transform [26], Riesz transforms [27], the martingale transform [32], the square function [12], [24] and well localized dyadic operators [15], [17], [7]. It is now also known for Calderón-Zygmund convolutions operators that are smooth averages of well localized operators [31].

In this paper, we are interested in obtaining sharp weight inequalities for the commutators of the Hilbert, Riesz, and Beurling transforms with multiplication by a locally integrable function $b \in BMO$.

1.1 Commutators: main results

Commutator operators are widely encountered and studied in many problems in PDEs, and Harmonic Analysis. One classical result of Coifman, Rochberg, and Weiss states in [6] that, for the Calderón-Zygmund singular integral operator with a smooth kernel, $[b,T]f := bT(f) - T(bf)$ is a bounded operator on $L^p_{\mathbb{R}^n}$, $1 < p < \infty$, when b is a BMO function. Weighted estimates for the commutator have been studied in [1], [20], [21], and [23]. Note that the commutator $[b,T]$ is more singular than the associated singular integral operator T, in particular, it does not satisfy the corresponding weak $(1,1)$ estimate. However one can find a weaker estimate in [21]. In 1997, C. Pérez [21] obtained the following result concerning commutators of singular integrals, for $1 < p < \infty$,

$$\|[b,T]f\|_{L^p(w)} \leq C\|b\|_{BMO}[w]^2_{A_\infty}\|M^2 f\|_{L^p(w)}\,,$$

where $M^2 = M \circ M$ denotes the Hardy-Littlewood maximal function iterated twice. With this result and Buckley's sharp estimate for the maximal function [3] one can immediately conclude that

$$\|[b,T]\|_{L^p(w)\to L^p(w)} \leq C[w]^2_{A_\infty}[w]^{\frac{2}{p-1}}_{A_p}\|b\|_{BMO}\,.$$

In this paper we show that for T the Hilbert, Riesz, Beurling transform, for $1 < p \leq 2$ one can drop $[w]_{A_\infty}$ term, in the above estimate, and this is sharp. However for $p > 2$, the $L^p(w)$-norm of $[b,T]$ is bounded above by $\|b\|_{BMO^d}[w]^2_{A^2_p}$. For $T = H$ the Hilbert transform we prove, using Bellman function techniques similar to those used in [2], [26], the following Theorem.

Theorem 1.1. *There exists a constant* $C > 0$*, such that*

$$\|[b,H]f\|_{L^p(w)\to L^p(w)} \leq C\|b\|_{BMO}[w]_{A_p}^{2\max\{1,\frac{1}{p-1}\}}\|f\|_{L^p(w)}\,,$$

and this is sharp for $1 < p < \infty$*.*

Most of the work goes into showing the quadratic estimate for $p = 2$, sharp extrapolation [8] then provides the right rate of growth for $p \neq 2$. An example of C. Pérez shows the rates are sharp. Our method involves the use of the dyadic paraproduct π_b and its adjoint π_b^*, both of which obey linear estimates in $L^2(w)$, see [2], like the Hilbert transform. It also uses Petermichl description of the Hilbert transform as an average of dyadic shift operators S, [25] and reduces the estimate to obtaining corresponding estimates for the commutator $[b,S]$. After we decompose this commutator in three parts:

$$[b,S]f = [\pi_b,S]f + [\pi_b^*,S]f + [\lambda_b,S]f\,,$$

we estimate each commutator separately. This decomposition has been used before to analyze the commutator, [25], [15], [16]. For precise definitions and detail derivations, see Section 1.2. The first two commutators immediately give the desired quadratic estimates in $L^2(w)$ from the known linear bounds of the operators commuted. For the third commutator we can prove a better than quadratic bound, in fact a linear bound. The following Theorem will be the crucial part of the proof.

Theorem 1.2. *There exists a constant* $C > 0$*, such that*

$$\|[\lambda_b,S]\|_{L^2(w)\to L^2(w)} \leq C[w]_{A_2^d}\|b\|_{BMO^d}\,, \tag{1.2}$$

for all $b \in BMO^d$ *and* $w \in A_2^d$*.*

This theorem is an immediate consequence of results in [14], [17] and [7], since the operator $[\lambda_b,S]$ belongs to the class of Haar shift operators for which they can prove linear bounds. We present a different proof of this result and others, using Bellman function techniques and bilinear Carleson embedding theorems, very much in the spirit of [25] and [2]. These arguments were found independently by the author, and we think they can be of interest.

We then observe that for any Haar shift operator T as defined in [17] the commutator $[\lambda_b,T]$ is again a Haar shift operator, and therefore it obeys linear bounds in A_2-characteristic of the weight as in Theorem 1.2. As a consequence, we obtain quadratic bounds for all commutators of Haar shift operators and BMO function b. In particular, this holds true for Haar shift operators in $\mathbb{R}^n$ whose averages recover the Riesz transforms [27] and for martingale transforms in $\mathbb{R}^2$ whose averages recover the Beurling-Ahlfors operator [30], [9]. Extrapolation will provide $L^p(w)$ bounds which turn out to be sharp for the Riesz transforms and Beurling-Ahlfors operators as well. The following Theorem holds

Theorem 1.3. *Let* T_τ *be the first class of Haar shift operators of index* τ*. Its convex hull include the Hilbert transform, Riesz transforms, the Beurling-Ahlfors operator and so on. Then, there exists a constant* $C(\tau,n,p)$ *which only depend on* τ*,* n *and* p *such that*

$$\|[b,T_\tau]\|_{L^p(w)\to L^p(w)} \leq C(\tau,n,p)[w]_{A_p}^{2\max\{1,\frac{1}{p-1}\}}\|b\|_{BMO}$$

See Section 10 for definitions and precise statements of these results.

1.2 The Hilbert transform case

The bilinear operator

$$fH(g) + H(f)g$$

maps $L^2 \times L^2$ into H^1, here H is the Hilbert transform and H^1 is the real Hardy space defined by

$$H^1(\mathbb{R}) := \{f \in L^1(\mathbb{R}) :\ Hf \in L^1(\mathbb{R})\}$$

with norm

$$\|f\|_{H^1} = \|f\|_{L^1} + \|Hf\|_{L^1} .$$

Because the dual of H^1 is BMO, we will pair with a BMO function b. Using that $H^* = -H$, we obtain that

$$\langle\, fH(g) + H(f)g,\, b\,\rangle = \langle\, f,\, H(g)b - H(gb)\,\rangle .$$

Hence the operator $g \mapsto H(g)b - H(gb)$ should be L^2 bounded. This expression $H(g)b - H(gb)$ is called the commutator of H with the BMO function b. More generally, we define as follows.

Definition 1.4. The commutator of the Hilbert transform H with a function b is defined as

$$[b, H](f) = bH(f) - H(bf) .$$

Our main concern in this paper is to prove that the commutator $[b, H]$, as an operator from $L^2_{\mathbb{R}}(w)$ into $L^2_{\mathbb{R}}(w)$ is bounded by the square of the A_2-characteristic, $[w]_{A_2}$, of the weight times the BMO norm, of b, $\|b\|_{BMO}$, where

$$\|b\|_{BMO} := \sup_I \frac{1}{|I|} \int_I |\, b(x) - \langle b\rangle_I| dx .$$

The supremum is taken over all intervals in $\mathbb{R}$. Note that when we restrict the supremum to dyadic intervals this will define BMO^d and we denote this dyadic BMO norm by $\|\cdot\|_{BMO^d}$. We now state our main results.

Theorem 1.5. *There exists C such that for all $w \in A_2$,*

$$\|[b, H]\|_{L^2(w)\to L^2(w)} \le C[w]^2_{A_2}\|b\|_{BMO} ,$$

for all $b \in BMO$.

Once we have boundedness and sharpness for the crucial case $p = 2$, we can carry out the power of the A_p-characteristic, for any $1 < p < \infty$, using the sharp extrapolation theorem [8] to obtain Theorem 1.1. Furthermore, an example of C. Pérez [22] shows this quadratic power is sharp. In [25], S. Petermichl showed that the norm of the commutator of the Hilbert transform is bounded by the supremum of the norms of the commutator of certain shift operators. This result follows after writing the kernel of the Hilbert transform as a well chosen averages of certain dyadic shift operators discovered by Petermichl. More precisely, S. Petermichl showed there is a non zero constant C such that

$$\|[b, H]\| \le C \sup_{\alpha,r} \|[b, S^{\alpha,r}]\| , \tag{1.3}$$

where the dyadic shift operator $S^{\alpha,r}$ is defined by

$$S^{\alpha,r} f = \sum_{I\in\mathcal{D}^{\alpha,r}} \langle f, h_I\rangle(h_{I_-} - h_{I_+}) .$$

Where h_I denotes the Haar function associated to the interval I, $I_\pm$ denote the left and right halves of I, and $\mathcal{D}^{\alpha,r}$ is a shifted and dilated system of dyadic intervals. Denote by S the shift operator corresponding to the standard dyadic intervals $\mathcal{D}$. See the next section for a precise definition.

Let us consider a compactly supported $b \in BMO^d$ and $f \in L^2$. Expanding b and f in the Haar system associated to the dyadic intervals $\mathcal{D}$,

$$b(x) = \sum_{I\in\mathcal{D}} \langle b, h_I\rangle h_I(x), \quad f(x) = \sum_{J\in\mathcal{D}} \langle f, h_J\rangle h_J(x)\,;$$

formally, we get the multiplication of b and f to be broken into three terms,

$$bf = \pi_b^*(f) + \pi_b(f) + \lambda_b(f)\,, \tag{1.4}$$

where π_b is the dyadic paraproduct, π_b^* is its adjoint and $\lambda_b(\cdot) = \pi_{(\cdot)}b$, defined as follows

$$\pi_b^*(f)(x) := \sum_{I\in\mathcal{D}} \langle b, h_I\rangle \langle f, h_I\rangle h_I^2(x)\,,$$

$$\pi_b(f)(x) := \sum_{I\in\mathcal{D}} \langle b, h_I\rangle \langle f\rangle_I h_I(x)\,,$$

$$\lambda_b(f)(x) := \sum_{I\in\mathcal{D}} \langle b\rangle_I \langle f, h_I\rangle h_I(x)\,.$$

Thus, we have

$$[b, S] = [\pi_b^*, S] + [\pi_b, S] + [\lambda_b, S]\,, \tag{1.5}$$

where

$$S(f) = \sum_{I\in\mathcal{D}} \langle f, h_I\rangle (h_{I_-} - h_{I_+})$$

and we can estimate each term separately. Notice that both π_b and π_b^* are bounded operators in L^p for $b \in BMO$, despite the fact that multipication by b is not a bounded operator in L^2 unless b is bounded (L^∞). Therefore, λ_b can not be a bounded operator in L^p. However $[\lambda_b, S]$ will be bounded on $L^p(w)$ and will be better behaved than $[b, S]$. Decomposition (1.5) was used to analyze the commutator with the shift operator first by Petermichl in [25], but also Lacey in [15] and authors in [16] to analyze the iterated commutators. Since all estimates are independent on the dyadic grid, through out this paper we only deal with the dyadic shift operator S associated to the standard dyadic grid $\mathcal{D}$. For a single shift operator the hypothesis required on b and w are that they belong to dyadic BMO^d and A_2^d with respect to the underlying dyadic grid defining the operator. However since ultimately we want to average over all grids, we will need b and w belonging to BMO^d and A_2^d for all shifted and scaled dyadic grids, that we will have if $b \in BMO$ and $w \in A_2$, non-dyadic BMO and A_2. Beznosova has proved linear bounds for π_b and π_b^*, [2], together with Petermichl's [26] linear bounds for S, this immediately provides the quadratic bounds for $[\pi_b, S]$ and $[\pi_b^*, S]$. Theorem 1.5 will be proved once we show the quadratic estimate holds for $[\lambda_b, S]$. We can actually obtain a better linear estimate as in Theorem 1.2. Some terms in (1.5) do also obey linear bounds.

Theorem 1.6. *There exists C such that*

$$\|\pi_b^* S\|_{L^2(w)} + \|S\pi_b\|_{L^2(w)} \le C[w]_{A_2^d} \|b\|_{BMO^d}\,.$$

for all $b \in BMO^d$.

Note the three operators $[\lambda_b, S]$, $\pi_b^* S$ and $S\pi_b$ are generalized Haar shift operators for which there are now two different proofs of linear bounds on $L^2(w)$ with respect to $[w]_{A_2^d}$, [17] and [7], and in this paper we present a third proof. We are now ready to explain the organization of this paper. In Section 2 we will introduce notation and discuss some useful results about weighted Haar systems. In Section 3 we will start our discussion on how to find the linear bound for the term $[\lambda_b, S]$, most of which will be very similar to calculations performed in [26]. In Section 4 we will introduce a number of Lemmas and Theorems that will be used. In Section 5 we will finish the linear estimate for the term $[\lambda_b, S]$, and prove Theorem 1.5. In Section 6 we prove the linear bound for $\pi_b^* S$. In Section 7 we reduce the proof of the linear bound for $S\pi_b$ to verifying three embedding conditions, two are proved in this section, the third is proved in Section 8 using a Bellman function argument. In Section 9 we present the $L^p(w)$ estimate of the commutator with a Haar shift operator, Theorem 1.3. Finally, in Section 10 we provide the sharpness for the commutators of Hilbert, Riesz transforms and Beurling-Ahlfors operators.

2 Preliminaries and Notation

Let us now introduce the notation which will be used frequently through this paper. Even though the A_p conditions have already been introduced in (1.1), we will state the special case of this condition when $p = 2$, namely A_2^d since we will refer repeatedly to this. We say w belongs to A_2^d class, if

$$[w]_{A_2^d} := \sup_{I\in\mathcal{D}} \langle w\rangle_I \langle w^{-1}\rangle_I < \infty\,. \tag{2.1}$$

Here we take the supremum over all dyadic interval in $\mathbb{R}$. Note that if $w \in A_2$ then $w \in A_2^d$ and $[w]_{A_2^d} \le [w]_{A_2}$. Intervals of the form $[k2^{-j}, (k+1)2^{-j})$ for integers j, k are called dyadic intervals. Again, let us denote $\mathcal{D}$ the collection of all dyadic intervals, and let us denote $\mathcal{D}(J)$ the collection of all dyadic subintervals of J. For any interval $I \in \mathcal{D}$, there is a Haar function defined by

$$h_I(x) = \frac{1}{|I|^{1/2}}(\chi_{I_+}(x) - \chi_{I_-}(x))\,,$$

where χ_I denotes the characteristic function of the interval I, $\chi_I(x) = 1$ if $x \in I$, $\chi_I(x) = 0$ otherwise. It is a well known fact that the Haar system $\{h_I\}_{I\in\mathcal{D}}$ is an orthonormal system in $L^2_{\mathbb{R}}$. We also consider the different grids of dyadic intervals parametrized by α, r, defined by

$$\mathcal{D}^{\alpha,r} = \{\alpha + rI : I \in \mathcal{D}\}\,,$$

for $\alpha \in \mathbb{R}$ and positive r. For each grid $\mathcal{D}^{\alpha,r}$ of dyadic intervals, there are corresponding Haar functions $h_I^{\alpha,r}$, $I \in \mathcal{D}^{\alpha,r}$ that are an orthonormal system in $L^2_{\mathbb{R}}$. Let us introduce a proper orthonormal system for $L^2_{\mathbb{R}}(w)$ defined by

$$h_I^w := \frac{1}{w(I)^{1/2}}\left[\frac{w(I_-)^{1/2}}{w(I_+)^{1/2}}\chi_{I_+} - \frac{w(I_+)^{1/2}}{w(I_-)^{1/2}}\chi_{I_-}\right],$$

where $w(I) = \int_I w$. We define the weighted inner product by $\langle f, g\rangle_w = \int_{\mathbb{R}} fgw$. Then, every function $f \in L^2(w)$ can be written as

$$f = \sum_{I\in\mathcal{D}} \langle f, h_I^w\rangle_w h_I^w\,,$$

where the sum converges a.e. in $L^2(w)$. Moreover,

$$\|f\|^2_{L^2(w)} = \sum_{I\in\mathcal{D}} |\langle f, h_I^w\rangle_w|^2\,. \tag{2.2}$$

Again $\mathcal{D}$ can be replaced by $\mathcal{D}^{\alpha,r}$ and the corresponding weighted Haar functions are an orthonormal system in $L^2(w)$. For convenience we will observe basic properties of the disbalanced Haar system. First observe that $\langle h_K, h_I^w\rangle_w$ could be non-zero only if $I \supseteq K$, moreover, for any $I \supseteq K$,

$$|\langle h_K, h_I^w\rangle_w| \le \langle w\rangle_K^{1/2}\,. \tag{2.3}$$

Here is the the calculation that provides (2.3),

$$\begin{aligned}|\langle h_K, h_I^w\rangle_w| &= \left|\int \frac{1}{|K|^{1/2}w(I)^{1/2}}(\chi_{K_+}(x)-\chi_{K_-}(x))\left[\frac{w(I_-)^{1/2}}{w(I_+)^{1/2}}\chi_{I_+}(x) - \frac{w(I_+)^{1/2}}{w(I_-)^{1/2}}\chi_{I_-}(x)\right]w(x)dx\right| \\ &\le \frac{1}{|K|^{1/2}w(I)^{1/2}}\underbrace{\int_K \left|\frac{w(I_-)^{1/2}}{w(I_+)^{1/2}}\chi_{I_+}(x) + \frac{w(I_+)^{1/2}}{w(I_-)^{1/2}}\chi_{I_-}(x)\right| w(x)dx}_{A}\,.\end{aligned}$$

If $K \subset I_+$, then $A \le w(I_-)^{1/2}w(I_+)^{-1/2}w(K)$. Thus

$$|\langle h_K, h_I^w\rangle_w| \le \langle w\rangle_K^{1/2}\sqrt{\frac{w(K)w(I_-)}{w(I_+)w(I)}} \le \langle w\rangle_K^{1/2}\,.$$

Similarly, if $K \subset I_-$. If $K = I$, then $A \le 2\,w(K_-)^{1/2}w(K_+)^{1/2}$. Thus

$$|\langle h_K, h_I^w\rangle_w| = |\langle h_K, h_K^w\rangle_w| \le \langle w\rangle_K^{1/2}\,\frac{2\sqrt{w(K_-)w(K_+)}}{w(K)} \le \langle w\rangle_K^{1/2}\,.$$

Estimate (2.3) implies that $|\langle h_{\hat J}, h_{\hat J}^{w^{-1}}\rangle_{w^{-1}}\langle h_J, h_J^w\rangle_w| \le \sqrt{2}[w]_{A_2^d}^{1/2}$, where $\hat J$ is the parent of J,

$$\begin{aligned}|\langle h_{\hat J}, h_{\hat J}^{w^{-1}}\rangle_{w^{-1}}\langle h_J, h_J^w\rangle_w| &\le \langle w^{-1}\rangle_{\hat J}^{1/2}\langle w\rangle_J^{1/2} = \langle w^{-1}\rangle_{\hat J}^{1/2}\left(\frac{1}{|J|}\int_J w\right)^{1/2} \\ &= \langle w^{-1}\rangle_{\hat J}^{1/2}\left(\frac{2}{|\hat J|}\int_J w\right)^{1/2} \le \sqrt{2}\langle w^{-1}\rangle_{\hat J}^{1/2}\langle w\rangle_{\hat J}^{1/2} \\ &\le \sqrt{2}\,[w]_{A_2^d}^{1/2}\,. \end{aligned}\tag{2.4}$$

Also, one can deduce similarly the following estimate

$$|\langle h_J, h_J^{w^{-1}}\rangle_{w^{-1}}\langle h_{J_-}, h_J^w\rangle_w| \le \sqrt{2}[w]_{A_2^d}^{1/2}\,. \tag{2.5}$$

For $I \supsetneq J$, h_I^w is constant on J. We will denote this constant by $h_I^w(J)$. Then $h_{\hat J}^w(J)$ is the constant value of $h_{\hat J}^w$ on J and $|h_{\hat J}^w(J)| \le w(J)^{-1/2}$, as can be seen by the following estimate,

$$|h_{\hat J}^w(J)| = \begin{cases} w(\hat J_-)^{1/2}/w(\hat J)^{1/2}w(\hat J_+)^{1/2} \le w(J)^{-1/2} & \text{if } J = \hat J_+ \\ w(\hat J_+)^{1/2}/w(\hat J)^{1/2}w(\hat J_-)^{1/2} \le w(J)^{-1/2} & \text{if } J = \hat J_-\,. \end{cases} \tag{2.6}$$

Let us define the weighted averages, $\langle g\rangle_{J,w} := w(J)^{-1}\int_J g(x)w(x)dx$. As with the standard Haar system, we can write the weighted averages

$$\langle g\rangle_{J,w} = \sum_{I\in\mathcal{D}: I\supsetneq J} \langle g, h_I^w\rangle_w h_I^w(J)\,. \tag{2.7}$$

In fact, here is the derivation of (2.7).

$$
\begin{aligned}
\langle g\rangle_{J,w} &= \frac{1}{w(J)}\int_J \sum_{I\in\mathcal{D}}\langle g, h_I^w\rangle_w h_I^w(x)w(x)dx = \frac{1}{w(J)}\int_J \sum_{I\in\mathcal{D}}\langle g, h_I^w\rangle_w h_I^w(J)w(x)dx \\
&= \frac{1}{w(J)}\int_J w(x)dx \sum_{I\in\mathcal{D}:I\supsetneq J}\langle g, h_I^w\rangle_w h_I^w(J) = \sum_{I\in\mathcal{D}:I\supsetneq J}\langle g, h_I^w\rangle_w h_I^w(J)\,.
\end{aligned}
$$

Also, we will be using system of functions $\{H_I^w\}_{I\in\mathcal{D}}$ defined by

$$
H_I^w = h_I\sqrt{|I|} - A_I^w\chi_I \quad \text{where} \quad A_I^w = \frac{\langle w\rangle_{I_+} - \langle w\rangle_{I_-}}{2\langle w\rangle_I}\,. \tag{2.8}
$$

Then, $\{w^{1/2}H_I^w\}$ is orthogonal in L^2 with norms satisfying the inequality $\|w^{1/2}H_I^w\|_{L^2} \le \sqrt{|I|\langle w\rangle_I}$, refer to [2]. Moreover, by Bessel's inequality we have, for all $g\in L^2$,

$$
\sum_{I\in\mathcal{D}}\frac{1}{|I|\langle w\rangle_I}\langle g, w^{1/2}H_I^w\rangle^2 \le \|g\|_{L^2}^2\,. \tag{2.9}
$$

3 Linear bound for $[\lambda_b, S]$ part I

In general, when we analyse commutator operators, a subtle cancelation delivers the result one wants to find. In the analysis of the commutator $[b,S]$, the part $[\lambda_b, S]$ will allow for certain cancelation. First, let us rewrite $[\lambda_b, S]$.

$$
\begin{aligned}
[\lambda_b, S](f) &= \lambda_b(Sf) - S(\lambda_b f) \\
&= \sum_{I\in\mathcal{D}}\langle b\rangle_I\langle Sf, h_I\rangle h_I - \sum_{J\in\mathcal{D}}\langle \lambda_b f, h_J\rangle(h_{J_-} - h_{J_+}) \\
&= \sum_{I\in\mathcal{D}}\sum_{J\in\mathcal{D}}\langle b\rangle_I\langle f, h_J\rangle\langle h_{J_-} - h_{J_+}, h_I\rangle h_I - \sum_{J\in\mathcal{D}}\sum_{I\in\mathcal{D}}\langle b\rangle_I\langle f, h_I\rangle\langle h_I, h_J\rangle(h_{J_-} - h_{J_+}) \\
&= \sum_{J\in\mathcal{D}}\langle b\rangle_{J_-}\langle f, h_J\rangle h_{J_-} - \sum_{J\in\mathcal{D}}\langle b\rangle_{J_+}\langle f, h_J\rangle h_{J_+} - \sum_{J\in\mathcal{D}}\frac{\langle b\rangle_{J_+} + \langle b\rangle_{J_-}}{2}\langle f, h_J\rangle(h_{J_-} - h_{J_+}) \\
&= \sum_{J\in\mathcal{D}}\frac{\langle b\rangle_{J_-} - \langle b\rangle_{J_+}}{2}\langle f, h_J\rangle h_{J_-} - \sum_{J\in\mathcal{D}}\frac{\langle b\rangle_{J_+} - \langle b\rangle_{J_-}}{2}\langle f, h_J\rangle h_{J_+} \\
&= -\sum_{J\in\mathcal{D}}\Delta_J b\langle f, h_J\rangle(h_{J_+} + h_{J_-})\,,
\end{aligned}
$$

recall the notation $\Delta_J\, b = (\langle b\rangle_{J_+} - \langle b\rangle_{J_-})/2$. To find the $L^2(w)$ operator norm of $[\lambda_b, S]$, it is enough to deal with the operator

$$
S_b(f) = \sum_{I\in\mathcal{D}}\Delta_I\, b\langle f, h_I\rangle h_{I_-}\,.
$$

We shall state the weighted operator norm of S_b as a Theorem and give a detailed proof. Theorem 1.2 is a direct consequence of the following Theorem. We will prove the following Theorem by the technique used in [26].

Theorem 3.1. *There exists a constant $C>0$, such that*

$$
\|S_b\|_{L^2(w)\to L^2(w)} \le C[w]_{A_2^d}\|b\|_{BMO^d} \tag{3.1}
$$

for all $b\in BMO^d$ and $w\in A_2^d$ for all $f\in L^2(w)$.

One can easily check that choosing $b = |I|^{1/2} h_I$, yields $\|S\|_{L^2(w)\to L^2(w)} \le C[w]_{A_2^d}$. Inequality (3.1) is equivalent to the following inequality for any positive function f, g

$$|\langle S_{b,w^{-1}} f, g\rangle_w| \le C[w]_{A_2^d} \|b\|_{BMO^d} \|f\|_{L^2(w^{-1})} \|g\|_{L^2(w)} , \tag{3.2}$$

where $S_{b,w^{-1}}(f) = S_b(w^{-1}f)$. Expanding f and g in the disbalanced Haar systems respectively for $L^2(w^{-1})$ and $L^2(w)$ yields for (3.2),

$$\begin{aligned} |\langle S_{b,w^{-1}} f, g\rangle_w| &= \Big| \int S_{b,w^{-1}} \big(\sum_{I\in\mathcal{D}} \langle f, h_I^{w^{-1}}\rangle_{w^{-1}} h_I^{w^{-1}} \big) \big(\sum_{J\in\mathcal{D}} \langle g, h_J^w\rangle_w h_J^w \big) w dx \Big| \\ &= \Big| \sum_{I\in\mathcal{D}} \sum_{J\in\mathcal{D}} \langle f, h_I^{w^{-1}}\rangle_{w^{-1}} \langle g, h_J^w\rangle_w \int h_J^w S_{b,w^{-1}}(h_I^{w^{-1}}) w dx \Big| \\ &= \Big| \sum_{I\in\mathcal{D}} \sum_{J\in\mathcal{D}} \langle f, h_I^{w^{-1}}\rangle_{w^{-1}} \langle g, h_J^w\rangle_w \langle S_{b,w^{-1}} h_I^{w^{-1}}, h_J^w\rangle_w \Big| . \end{aligned} \tag{3.3}$$

In (3.3),

$$\langle S_{b,w^{-1}} h_I^{w^{-1}}, h_J^w\rangle_w = \Big\langle \sum_{L\in\mathcal{D}} \Delta_L b \langle h_L, w^{-1} h_I^{w^{-1}} \rangle \, h_{L^-}, h_J^w \Big\rangle_w = \sum_{L\in\mathcal{D}} \Delta_L b \langle h_L, h_I^{w^{-1}}\rangle_{w^{-1}} \langle h_{L^-}, h_J^w\rangle_w$$

Since $\langle h_L, h_I^{w^{-1}}\rangle_{w^{-1}} \neq 0$, only when $L \subseteq I$ and $\langle h_{L^-}, h_J^w\rangle_w \neq 0$ only when $L_- \subseteq J$, then we have non-zero terms if $I \subseteq J$ or $\hat{J} \subseteq I$ in the sum of (3.3). Thus we can split the sum into four parts, $\sum_{I=J}$, $\sum_{I=\hat{J}}$, $\sum_{\hat{J}\subsetneq I}$, and $\sum_{I\subsetneq J}$. Let us now introduce the truncated shift operator

$$S_b^I(f) := \sum_{L\in\mathcal{D}(I)} \Delta_L b \langle f, h_L\rangle h_{L_-} ,$$

and its composition with multiplication by w^{-1},

$$S_{b,w^{-1}}^I(f) := \sum_{L\in\mathcal{D}(I)} \Delta_L b \langle w^{-1} f, h_L\rangle h_{L_-} .$$

We will see that the weighted norm $\|S_{b,w^{-1}}^I \chi_I\|_{L^2(w)}$, plays a main role in our estimate for $\langle S_{b,w^{-1}} h_I^{w^{-1}}, h_J^w\rangle_w$.

4 Theorems and Lemmas

To prove inequality (3.2) we need several theorems and lemmas. One can find the proof in the indicated references. First we recall some embedding theorems. Another main result in [26] is a two-weighted bilinear embedding theorem, which was proved by a Bellman function argument.

Theorem 4.1 (Petermichl's Bilinear Embedding Theorem). *Let w and v be weights so that $\langle w\rangle_I \langle v\rangle_I \le Q$ for all intervals I and let $\{\alpha_I\}$ be a non-negative sequence so that the three estimates below hold for all J*

$$\sum_{I\in\mathcal{D}(J)} \frac{\alpha_I}{\langle w\rangle_I} \le Q\, v(J) \tag{4.1}$$

$$\sum_{I\in\mathcal{D}(J)} \frac{\alpha_I}{\langle v\rangle_I} \le Q\, w(J) \tag{4.2}$$

$$\sum_{I\in\mathcal{D}(J)} \alpha_I \le Q\,|J|\,. \tag{4.3}$$

Then there is c such that for all $f \in L^2(w)$ and $g \in L^2(v)$

$$\sum_{I\in\mathcal{D}} \alpha_I \langle f\rangle_{I,w}\langle g\rangle_{I,v} \le cQ\|f\|_{L^2(w)}\|g\|_{L^2(v)}\,.$$

Corollary 4.2 (Bilinear Embedding Theorem)**.** *Let w and v be weights. Let $\{\alpha_I\}$ be a sequence of nonnegative numbers such that for all dyadic intervals $J \in \mathcal{D}$ the following three inequalities hold with some constant $Q > 0$,*

$$\sum_{I\in\mathcal{D}(J)} \alpha_I\langle v\rangle_I\,|I| \le Q\,v(J) \tag{4.4}$$

$$\sum_{I\in\mathcal{D}(J)} \alpha_I\langle w\rangle_I\,|I| \le Q\,w(J) \tag{4.5}$$

$$\sum_{I\in\mathcal{D}(J)} \alpha_I\langle w\rangle_I\langle v\rangle_I \le Q\,|J|\,. \tag{4.6}$$

Then for any two nonnegative function $f,\, g \in L^2$

$$\sum_{I\in\mathcal{D}} \alpha_I\langle fv^{1/2}\rangle_I\langle gw^{1/2}\rangle_I\,|I| \le CQ\|f\|_{L^2}\|g\|_{L^2}$$

holds with some constant $C > 0$.

Both Bilinear Embedding Theorems are key tools in our estimate. One version of such a theorem appeared in [19]. The original version of the next lemma also appeared in [26]. Using the fact that, for all $I \in \mathcal{D}$, $|\Delta_I b| \le 2\|b\|_{BMO^d}$, one can prove the following lemma similarly to its original proof.

Lemma 4.3. *There is a constant c such that $\|S^I_{b,w^{-1}}\chi_I\|_{L^2(w)} \le c\|b\|_{BMO^d}[w]_{A_2^d}w^{-1}(I)^{1/2}$ for all intervals I and weights $w \in A_2^d$.*

We will also need the Weighted Carleson Embedding Theorem from [19], and some other inequalities for weights.

Theorem 4.4 (Weighted Carleson Embedding Theorem)**.** *Let $\{\alpha_J\}$ be a non-negative sequence such that for all dyadic intervals I*

$$\sum_{J\in\mathcal{D}(I)} \alpha_J \le Qw^{-1}(I)\,.$$

Then for all $f \in L^2(w^{-1})$

$$\sum_{J\in\mathcal{D}} \alpha_J\langle f\rangle^2_{J,w^{-1}} \le 4Q\|f\|^2_{L^2(w^{-1})}\,.$$

Theorem 4.5 (Wittwer's sharp version of Buckley's inequality)**.** *There exist a positive constant C such that for any weight $w \in A_2^d$ and dyadic interval $I \in \mathcal{D}$,*

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)} \frac{\big(\langle w\rangle_{I_+} - \langle w\rangle_{I_-}\big)^2}{\langle w\rangle_I}\,|I| \le C[w]_{A_2^d}\langle w\rangle_J\,.$$

We refer to [32] for the proof. You can find extended versions of Theorem 4.4 and 4.5 to the doubling positive measure σ in [24]. One can find the Bellman function proof of the following three Lemmas in [2].

Lemma 4.6. *For all dyadic interval J and all weights w.*

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|I||\Delta_I w|^2\frac{1}{\langle w\rangle_I^3}\le\langle w^{-1}\rangle_J\,.$$

Lemma 4.7. *Let w be a weight and $\{\alpha_I\}$ be a Carleson sequence of nonnegative numbers. If there exist a constant $Q>0$ such that*

$$\forall\, J\in\mathcal{D}\,,\quad\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\alpha_I\le Q\,,$$

then

$$\forall\, J\in\mathcal{D}\,,\quad\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\frac{\alpha_I}{\langle w^{-1}\rangle_I}\le 4Q\langle w\rangle_J$$

and therefore if $w\in A_2^d$ the for any $J\in\mathcal{D}$ we have

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle w\rangle_I\alpha_I\le 4Q[w]_{A_2^d}\langle w\rangle_J\,.$$

Lemma 4.8. *If $w\in A_2^d$ then there exists a constant $C>0$ such that*

$$\forall J\in\mathcal{D}\,,\quad\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\left(\frac{\langle w\rangle_{I_+}-\langle w\rangle_{I_-}}{\langle w\rangle_I}\right)^2|\,I|\,\langle w\rangle_I\langle w^{-1}\rangle_I\le C[w]_{A_2^d}\,.$$

5 Linear bound for $[\lambda_b, S]$ part II

We will continue to estimate the sum (3.3) in four parts.

5.1 $\sum_{I=\hat{J}}$

For this case, it is sufficient to show that

$$|\langle S_{b,w^{-1}}h_{\hat{J}}^{w^{-1}},h_J^w\rangle_w|\le c\|b\|_{BMO^d}[w]_{A_2^d}\,.$$

Since $\langle h_k,h_I^w\rangle_w$ could be non-zero only if $K\subseteq I$,

$$|\langle S_{b,w^{-1}}h_{\hat{J}}^{w^{-1}},h_J^w\rangle_w|=\Big|\Big\langle\sum_{L\in\mathcal{D}}\Delta_L b\,\langle w^{-1}h_L^{w^{-1}},h_L\rangle h_{L_-},h_J^w\Big\rangle_w\Big|=\Big|\sum_{L\in\mathcal{D}}\Delta_L b\,\langle h_{\hat{J}}^{w^{-1}},h_L\rangle_{w^{-1}}\langle h_{L_-},h_J^w\rangle_w\Big|$$

has non-zero term only when $L\subseteq\hat{J}$. Thus

$$\begin{aligned}|\langle S_{b,w^{-1}}h_L^{w^{-1}},h_J^w\rangle_w|&=\Big|\sum_{L\in\mathcal{D}(\hat{J}\,)}\Delta_L b\,\langle h_{\hat{J}}^{w^{-1}},h_L\rangle_{w^{-1}}\langle h_{L_-},h_J^w\rangle_w\Big|\\&=\Big|\sum_{L\in\mathcal{D}(J)}\Delta_L b\,\langle h_{\hat{J}}^{w^{-1}},h_L\rangle_{w^{-1}}\langle h_{L_-},h_J^w\rangle_w\Big|+\Big|\sum_{L\in\mathcal{D}(J^s)}\Delta_L b\,\langle h_{\hat{J}}^{w^{-1}},h_L\rangle_{w^{-1}}\langle h_{L_-},h_J^w\rangle_w\Big|\\&\quad+|\Delta_{\hat{J}}\,b\,\langle h_{\hat{J}}^{w^{-1}},h_{\hat{J}}\rangle_{w^{-1}}\langle h_{\hat{J}_-},h_J^w\rangle_w|\\&\le|\langle S_{b,w^{-1}}^J h_{\hat{J}}^{w^{-1}},h_J^w\rangle_w|+|\Delta_{\hat{J}}\,b\,\langle h_{\hat{J}}^{w^{-1}},h_{\hat{J}}\rangle_{w^{-1}}\langle h_{\hat{J}_-},h_J^w\rangle_w|\end{aligned}$$

in the second equality, J^s denotes the sibling of J, so for all $L \subseteq J^s$, $\langle h_{L_-}, h_J^w\rangle_w = 0$. Then, by (2.4),

$$|\Delta_{\hat{J}}\, b\, \langle h_{\hat{J}}^{w^{-1}}, h_{\hat{J}}\rangle_{w^{-1}} \langle h_{\hat{J}_-}, h_J^w\rangle_w| \leq \sqrt{2}\|b\|_{BMO^d}[w]_{A_2^d}^{1/2}\,. \tag{5.1}$$

So for the remaining part:

$$|\langle S^J_{b,w^{-1}} h_{\hat{J}}^{w^{-1}}, h_J^w\rangle_w| = |h_{\hat{J}}^{w^{-1}}(J)\langle S^J_{b,w^{-1}}\chi_J, h_J^w\rangle_w| \leq c\|b\|_{BMO^d}[w]_{A_2^d}\,, \tag{5.2}$$

here the last inequality uses (2.6) and Lemma 4.3.

5.2 $\sum_{I=J}$

In this case, the argument is similar to the argument in Section 5.1. We have

$$|\langle S_{b,w^{-1}} h_J^{w^{-1}}, h_J^w\rangle_w| = |\sum_{L\in\mathcal{D}} \Delta_L b\, \langle h_J^{w^{-1}}, h_L\rangle_{w^{-1}} \langle h_{L_-}, h_J^w\rangle_w|$$

here we have zero summands, unless $L \subseteq J$. Thus,

$$\begin{aligned}|\langle S_{b,w^{-1}} h_J^{w^{-1}}, h_J^w\rangle_w| &= |\langle S^J_{b,w^{-1}} h_J^{w^{-1}}, h_J^w\rangle_w| \\ &\leq |\langle S^{J_+}_{b,w^{-1}} h_J^{w^{-1}}, h_J^w\rangle_w| + |\langle S^{J_-}_{b,w^{-1}} h_J^{w^{-1}}, h_J^w\rangle_w| + |\Delta_J\, b\, \langle h_J^{w^{-1}}, h_J\rangle_{w^{-1}} \langle h_{J_-}, h_J^w\rangle_w| \\ &\leq c\,\|b\|_{BMO^d}[w]_{A_2^d}\,.\end{aligned}$$

In the last inequality, we use same arguments as in (5.2) for the first two term, and (2.5) for the last term.

5.3 $\sum_{\hat{J}\subsetneq I}$ and $\sum_{I\subsetneq J}$

To obtain our desired results, we need to understand the supports of $S_b(w^{-1}h_I^{w^{-1}})$ and $S_b^*(wh_J^w)$. Since

$$S_b(w^{-1}h_I^{w^{-1}}) = \sum_{L\in\mathcal{D}} \Delta_L b\, \langle w^{-1}h_I^{w^{-1}}, h_L\rangle h_{L_-} = \sum_{L\in\mathcal{D}} \Delta_L b\, \langle h_I^{w^{-1}}, h_L\rangle_{w^{-1}} h_{L_-}\,,$$

and $\langle h_I^{w^{-1}}, h_L\rangle$ can be non-zero only when $L \subseteq I$, $S_b(w^{-1}h_I^{w^{-1}})$ is supported by I. Also,

$$\langle S_b(w^{-1}h_I^{w^{-1}}), h_J^w\rangle_w = \langle h_I^{w^{-1}}, S_b^*(wh_J^w)\rangle_{w^{-1}}\,, \tag{5.3}$$

yield that $S_b^*(wh_J^w) = \sum_{L\in\mathcal{D}} \Delta_L b\langle wh_J^w, h_{L_-}\rangle h_L$ is supported by $\hat{J}$. Let us now consider the sum $\hat{J} \subsetneq I$. Then

$$\begin{aligned}
\Big| \sum_{I,J:\hat{J}\subsetneq I} \langle f, h_I^{w^{-1}}\rangle_{w^{-1}} \langle g, h_J^w\rangle_w \langle S_{b,w^{-1}} h_I^{w^{-1}}, h_J^w\rangle_w \Big| \qquad & \\
= \Big| \sum_{I,J:\hat{J}\subsetneq I} \langle f, h_I^{w^{-1}}\rangle_{w^{-1}} \langle g, h_J^w\rangle_w \langle h_I^{w^{-1}}, S_b^*(wh_J^w)\rangle_{w^{-1}} \Big| & \\
= \Big| \sum_{J\in\mathcal{D}} \sum_{I:I\supsetneq\hat{J}} \langle f, h_I^{w^{-1}}\rangle_{w^{-1}} \langle g, h_J^w\rangle_w h_I^{w^{-1}}(\hat{J}) \langle S_{b,w^{-1}}\chi_{\hat{J}}, h_J^w\rangle_w \Big| & \qquad (5.4)\\
= \Big| \sum_{J\in\mathcal{D}} \langle f\rangle_{\hat{J},w^{-1}} \langle g, h_J^w\rangle_w \langle S_{b,w^{-1}}\chi_{\hat{J}}, h_J^w\rangle_w \Big| & \qquad (5.5)\\
\le \|g\|_{L^2(w)} \bigg(\sum_{J\in\mathcal{D}} \langle f\rangle_{\hat{J},w^{-1}}^2 \langle S_{b,w^{-1}}\chi_{\hat{J}}, h_J^w\rangle_w^2 \bigg)^{1/2}, & \qquad (5.6)
\end{aligned}$$

here (5.3) and the fact that $S_b^*(wh_J^w)$ is supported by $\hat{J}$ are used for equality (5.4), and (5.5) uses (2.7) and (5.3). If we show that

$$\sum_{J\in\mathcal{D}} \langle f\rangle_{\hat{J},w^{-1}}^2 \langle S_{b,w^{-1}}\chi_{\hat{J}}, h_J^w\rangle_w^2 \le c\|b\|_{BMO^d}^2 [w]_{A_2^d}^2 \|f\|_{L^2(w^{-1})}^2, \qquad (5.7)$$

then we have

$$\Big| \sum_{I,J:\hat{J}\subsetneq I} \langle f, h_I^{w^{-1}}\rangle_{w^{-1}} \langle g, h_J^w\rangle_w \langle S_{b,w^{-1}} h_I^{w^{-1}}, h_J^w\rangle_w \Big| \le C\|b\|_{BMO^d} [w]_{A_2^d} \|f\|_{L(w^{-1})}^2 \|g\|_{L^2(w)}.$$

To prove the inequality (5.7), we apply the Theorem 4.4. The embedding condition becomes

$$\sum_{J\in\mathcal{D}:J\subsetneq I} \langle S_{b,w^{-1}}\chi_{\hat{J}}, h_J^w\rangle_w^2 \le c\|b\|_{BMO^d}^2 [w]_{A_2^d}^2 w^{-1}(I)$$

after shifting the indices. Since $\langle h_{L_-}, h_J^w\rangle_w = 0$ unless $L \subseteq \hat{J}$, and we will sum over J such that $J \subsetneq I$, we can write

$$\begin{aligned}
\langle S_{b,w^{-1}}\chi_{\hat{J}}, h_J^w\rangle_w &= \sum_{L\in\mathcal{D}} \Delta_L b\langle w^{-1}\chi_{\hat{J}}, h_L\rangle \langle h_{L_-}, h_J^w\rangle_w = \sum_{L\in\mathcal{D}(\hat{J})} \Delta_L b\langle w^{-1}\chi_{\hat{J}}, h_L\rangle \langle h_{L_-}, h_J^w\rangle_w \\
&= \sum_{L\in\mathcal{D}(I)} \Delta_L b\langle w^{-1}\chi_I, h_L\rangle \langle h_{L_-}, h_J^w\rangle_w = \langle S_{b,w^{-1}}^I \chi_I, h_J^w\rangle_w.
\end{aligned}$$

Thus,

$$\sum_{J\in\mathcal{D}:J\subsetneq I} \langle S_{b,w^{-1}}\chi_{\hat{J}}, h_J^w\rangle_w^2 = \sum_{J\in\mathcal{D}:J\subsetneq I} \langle S_{b,w^{-1}}^I \chi_I, h_J^w\rangle_w^2 \le \|S_{b,w^{-1}}^I \chi_I\|_{L^2(w)}^2,$$

last inequality due to (2.2). By Lemma 4.3, the embedding condition holds. Hence we are done for the sum $\hat{J} \subsetneq I$. The part $\sum_{I\subsetneq J}$ is similar to $\sum_{\hat{J}\subsetneq I}$. One uses that $S_b(w^{-1}h_I^{w^{-1}})$ is supported by I and Theorem 4.4.

5.4 Proof of Theorem 1.5

We refer to [2] for following theorem.

Theorem 5.1. *The norm of dyadic paraproduct on the weighted Lebesgue space $L^2(w)$ is bounded from above by a constant multiple of the product of the A_2^d characteristic of the weight w and the BMO^d norm of b.*

To break $[b,S]$ into three parts, as in (1.5), we assumed that $b \in BMO^d$ is compactly supported. However, we need to replace such a b with a general BMO^d function. In order to pass from a compactly supported b to general $b \in BMO^d$, we need the following lemma which is suggested in [10].

Lemma 5.2. *Suppose $\phi \in BMO$. Let $\widetilde{I}$ be the interval concentric with I having length $|\widetilde{I}| = 3|I|$. Then there is $\psi \in BMO$ such that $\psi = \phi$ on I, $\psi = 0$ on $\mathbb{R} \setminus \widetilde{I}$ and $\|\psi\|_{BMO} \le c\|\phi\|_{BMO}$.*

Proof. Without loss of generality, we assume $\langle \phi \rangle_I = 0$. Write $I = \bigcup_{n=0}^{\infty} J_n$ where $dist(J_n, \partial I) = |J_n|$, as in following figure.



Then J_0 is the middle third of I. For $n > 0$, let K_n be the reflection of J_n across the nearest endpoint of I and set

$$\psi(x) = \begin{cases} \phi(x) & \text{if } x \in I \\ \langle \phi \rangle_{J_n} & \text{if } x \in K_n \\ 0 & \text{otherwise}. \end{cases}$$

This construction of ψ satisfies Lemma 5.2.

□

By Theorem 3.1, Corollary 1.2, Theorem 5.1, Lemma 5.2 and using the fact $\|\pi_b\| = \|\pi_b^*\|$, we can prove Theorem 1.5.

Proof of Theorem 1.5. For any compactly supported $b \in BMO$,

$$\begin{aligned} \|[b,H]\|_{L^2(w)\to L^2(w)} &\le C \sup_{\alpha,r} \|[b, S^{\alpha,r}]\|_{L^2(w)\to L^2(w)} \\ &\le C \sup_{\alpha,r} \big(\|[\pi_b, S^{\alpha,r}]\|_{L^2(w)\to L^2(w)} + \|[\pi_b^*, S^{\alpha,r}]\|_{L^2(w)\to L^2(w)} + \|[\lambda_b, S^{\alpha,r}]\|_{L^2(w)\to L^2(w)} \big) \\ &\le C\big(4\|\pi_b\|_{L^2(w)\to L^2(w)} \sup_{\alpha,r} \|S^{\alpha,r}\|_{L^2(w)\to L^2(w)} + C[w]_{A_2}\|b\|_{BMO}\big) \\ &\le C[w]_{A_2}^2 \|b\|_{BMO}. \end{aligned}$$

For fixed b, we consider the sequence of intervals $I_k = [-k,k]$ and the sequence of BMO functions b_k which are constructed as in Lemma 5.2. Then, there is a constant c, which does not depend on k, such that $\|b_k\|_{BMO} \le c\|b\|_{BMO}$. Furthermore, there is a uniform constant C such that

$$\|[b_k, H]\|_{L^2(w)\to L^2(w)} \le C[w]_{A_2}^2 \|b\|_{BMO}. \tag{5.8}$$

Therefore, for some subsequence of integers k_j and $f \in L^2(w)$, $[b_{k_j}, H](f)$ converges to $[b,H](f)$ almost everywhere. Letting $j \to \infty$ and using Fatou's lemma, we deduce that (5.8) holds for all $b \in BMO$. □

6 Linear bound for $\pi_b^* S$

It might be useful to know what is the adjoint operator of S. Let us define $\operatorname{sgn}(I) = \pm 1$ if $I = \hat{I}_{\mp}$. Then, for any function $f, g \in L^2$,

$$\begin{aligned}
\langle Sf, g\rangle &= \sum_{I\in\mathcal{D}}\sum_{J\in\mathcal{D}}\langle f, h_I\rangle\langle g, h_J\rangle\langle h_{I_-} - h_{I_+}, h_J\rangle \\
&= \sum_{I\in\mathcal{D}}\langle f, h_I\rangle\langle g, h_{I_-}\rangle - \sum_{I\in\mathcal{D}}\langle f, h_I\rangle\langle g, h_{I_+}\rangle \\
&= \sum_{I\in\mathcal{D}}\langle f, h_I\rangle\big(\operatorname{sgn}(I_-)\langle g, h_{I_-}\rangle + \operatorname{sgn}(I_+)\langle g, h_{I_+}\rangle\big) \\
&= \sum_{I\in\mathcal{D}}\langle f, h_{\hat{I}}\rangle\operatorname{sgn}(I)\langle g, h_I\rangle \\
&= \Big\langle f, \sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle g, h_I\rangle h_{\hat{I}}\Big\rangle = \langle f, S^* g\rangle\,.
\end{aligned}$$

Now, we see the adjoint operator of dyadic shift operator S is

$$S^* f(x) = \sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle f, h_I\rangle h_{\hat{I}}(x)\,.$$

The following lemma provides the bound we are looking for the term $\pi_b^* S$.

Lemma 6.1. *Let $w \in A_2^d$ and $b \in BMO^d$. Then, there exists C so that*

$$\|\pi_b^* S\|_{L^2(w)\to L^2(w)} \le C[w]_{A_2^d}\|b\|_{BMO^d}\,.$$

Proof. In order to prove Lemma 6.1 it is enough to show that for any positive square integrable function f, g

$$\langle \pi_b^* S(fw^{-1/2}), gw^{1/2}\rangle \le C[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2}\|g\|_{L^2}\,. \tag{6.1}$$

Using the system of functions $\{H_I^w\}_{I\in\mathcal{D}}$ defined in (2.8), we can rewrite the left hand side of (6.1)

$$\begin{aligned}
\langle \pi_b^* S(fw^{-1/2}), gw^{1/2}\rangle = \langle S(fw^{-1/2}), \pi_b(gw^{1/2})\rangle &= \sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I\langle b, h_I\rangle\langle S(fw^{-1/2}), h_I\rangle \\
&= \sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I\langle b, h_I\rangle\operatorname{sgn}(I)\langle fw^{-1/2}, h_{\hat{I}}\rangle \\
&= \sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle gw^{1/2}\rangle_I\langle b, h_I\rangle\langle fw^{-1/2}, H_{\hat{I}}^{w^{-1}}\rangle\frac{1}{\sqrt{|\hat{I}|}} \\
&\quad + \sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle gw^{1/2}\rangle_I\langle b, h_I\rangle\langle fw^{-1/2}, A_{\hat{I}}^{w^{-1}}\chi_{\hat{I}}\rangle\frac{1}{\sqrt{|\hat{I}|}}\,.
\end{aligned} \tag{6.2}$$

Our claim is that both sums in (6.2) are bounded by $[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2}\|g\|_{L^2}$, i.e.

$$\sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle gw^{1/2}\rangle_I\langle b, h_I\rangle\langle fw^{-1/2}, H_{\hat{I}}^{w^{-1}}\rangle\frac{1}{\sqrt{|\hat{I}|}} \le C[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2}\|g\|_{L^2} \tag{6.3}$$

and

$$\sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle gw^{1/2}\rangle_I\langle b, h_I\rangle\langle fw^{-1/2}, A_{\hat{I}}^{w^{-1}}\chi_{\hat{I}}\rangle\frac{1}{\sqrt{|\hat{I}|}} \le C[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2}\|g\|_{L^2}\,. \tag{6.4}$$

First let us verify the bound for (6.3). Using Cauchy-Schwarz inequality,

$$\begin{aligned}\sum_{I\in\mathcal{D}} &\operatorname{sgn}(I)\langle gw^{1/2}\rangle_I\langle b,h_I\rangle\langle fw^{-1/2},H_{\hat I}^{w^{-1}}\rangle\frac{1}{\sqrt{|\hat I|}}\\ &\leq\Big(\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I^2\langle b,h_I\rangle^2\langle w^{-1}\rangle_{\hat I}\Big)^{1/2}\Big(\sum_{I\in\mathcal{D}}\frac{1}{|\hat I|\langle w^{-1}\rangle_{\hat I}}\langle f,w^{-1/2}H_{\hat I}^{w^{-1}}\rangle^2\Big)^{1/2}\\ &\leq\|f\|_{L^2}\Big(\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I^2\langle b,h_I\rangle^2\langle w^{-1}\rangle_{\hat I}\Big)^{1/2}. \qquad (6.5)\end{aligned}$$

Thus, for (6.3), it is enough to show that

$$\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I^2\langle b,h_I\rangle^2\langle w^{-1}\rangle_{\hat I}\leq C[w]_{A_2^d}^2\|b\|_{BMO^d}^2\|g\|_{L^2}^2\,. \qquad (6.6)$$

It is clear that $2\langle w\rangle_{\hat I}\geq\langle w\rangle_I$, thus

$$\begin{aligned}\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I^2\langle b,h_I\rangle^2\langle w^{-1}\rangle_{\hat I}&=\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I^2\langle b,h_I\rangle^2\langle w^{-1}\rangle_{\hat I}\langle w\rangle_I\langle w\rangle_I^{-1}\\ &\leq 2[w]_{A_2^d}\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I^2\langle b,h_I\rangle^2\langle w\rangle_I^{-1}\,.\end{aligned}$$

If we show for all $J\in\mathcal{D}$,

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w\rangle_I^{-1}\langle w\rangle_I^2=\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w\rangle_I\leq[w]_{A_2^d}\|b\|_{BMO^d}^2\langle w\rangle_J\,, \qquad (6.7)$$

then by Weighted Carleson Embedding Theorem 4.4 with w instead of w^{-1}, we will have (6.6). Since $b\in BMO^d$, $\{\langle b,h_I\rangle^2\}_{I\in\mathcal{D}}$ is a Carleson sequence with constant $\|b\|_{BMO^d}^2$ that is

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\leq\|b\|_{BMO^d}^2\,.$$

Applying Lemma 4.7 with $\alpha_I=\langle b,h_I\rangle$ we have inequality (6.7). We now concentrate on the estimate (6.4), we can estimate the left hand side of (6.4) as follows.

$$\begin{aligned}\sum_{I\in\mathcal{D}} &\operatorname{sgn}(I)\langle gw^{1/2}\rangle_I\langle b,h_I\rangle\langle fw^{-1/2},A_{\hat I}^{w^{-1}}\chi_{\hat I}\rangle\frac{1}{\sqrt{|\hat I|}}\\ &=\sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle gw^{1/2}\rangle_I\langle b,h_I\rangle\langle fw^{-1/2}\rangle_{\hat I}A_{\hat I}^{w^{-1}}\sqrt{|\hat I|}\\ &\leq\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I\,|\langle b,h_I\rangle|\,\langle fw^{-1/2}\rangle_{\hat I}\,|A_{\hat I}^{w^{-1}}|\sqrt{|\hat I|}\\ &\leq 2\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_{\hat I}\,|\langle b,h_I\rangle|\,\langle fw^{-1/2}\rangle_{\hat I}\,|A_{\hat I}^{w^{-1}}|\sqrt{|\hat I|}\\ &=2\sum_{I\in\mathcal{D}}\langle gw^{1/2}\rangle_I\,\big(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|\big)\,\langle fw^{-1/2}\rangle_I\,|A_I^{w^{-1}}|\sqrt{|I|}\,.\end{aligned}$$

By Bilinear Embedding Theorem, inequality (6.4) holds provided the following three inequalities hold,

$$\forall\, J\in\mathcal{D}, \frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\big(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|\big)|A_I^{w^{-1}}|\sqrt{|I|}\langle w^{-1}\rangle_I\langle w\rangle_I\le C\|b\|_{BMO^d}\,[w^{-1}]_{A_2^d}\,, \tag{6.8}$$

$$\forall\, J\in\mathcal{D}, \frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\big(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|\big)|A_I^{w^{-1}}|\sqrt{|I|}\langle w^{-1}\rangle_I\le C\|b\|_{BMO^d}\,[w^{-1}]_{A_2^d}\langle w^{-1}\rangle_J\,, \tag{6.9}$$

$$\forall\, J\in\mathcal{D}, \frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\big(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|\big)|A_I^{w^{-1}}|\sqrt{|I|}\langle w\rangle_I\le C\|b\|_{BMO^d}\,[w^{-1}]_{A_2^d}\langle w\rangle_J\,. \tag{6.10}$$

For (6.8), by Cauchy-Schwarz inequality

$$\begin{aligned}&\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\big(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|\big)|A_I^{w^{-1}}|\sqrt{|I|}\langle w^{-1}\rangle_I\langle w\rangle_I\\ &\le\Bigg(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\big(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|\big)^2\langle w^{-1}\rangle_I\langle w\rangle_I\Bigg)^{1/2}\Bigg(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(A_I^{w^{-1}})^2|I|\langle w^{-1}\rangle_I\langle w\rangle_I\Bigg)^{1/2}.\end{aligned}$$

Since

$$\sum_{I\in\mathcal{D}}(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|)^2\le 3\sum_{I\in\mathcal{D}}\langle b,h_I\rangle^2\,, \tag{6.11}$$

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|)^2\langle w^{-1}\rangle_I\langle w\rangle_I\le C[w^{-1}]_{A_2^d}\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\le C[w^{-1}]_{A_2^d}\|b\|^2_{BMO^d}\,,$$

and by Lemma 4.8,

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(A_I^{w^{-1}})^2|I|\langle w^{-1}\rangle_I\langle w\rangle_I\le C\,[w^{-1}]_{A_2^d}\,.$$

Thus embedding condition (6.8) holds. For (6.9), by Cauchy-Schwarz inequality and (6.11) we have

$$\begin{aligned}&\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(|\langle b,h_{I_-}\rangle|+|\langle b,h_{I_+}\rangle|)\,|A_I^{w^{-1}}|\sqrt{|I|}\langle w^{-1}\rangle_I\\ &\qquad\le C\Bigg(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w^{-1}\rangle_I\Bigg)^{1/2}\Bigg(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(A_I^{w^{-1}})^2|I|\langle w^{-1}\rangle_I\Bigg)^{1/2}.\end{aligned}$$

By Theorem 4.5,

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(A_I^{w^{-1}})^2|I|\langle w^{-1}\rangle_I=\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\left(\frac{\langle w^{-1}\rangle_{I_+}-\langle w^{-1}\rangle_{I_-}}{2\langle w^{-1}\rangle_I}\right)^2|I|\langle w^{-1}\rangle_I\le C[w^{-1}]_{A_2^d}\langle w^{-1}\rangle_J\,.$$

Similarly with (6.7), we have

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w^{-1}\rangle_I\le[w^{-1}]_{A_2}\|b\|^2_{BMO^d}\langle w^{-1}\rangle_J\,.$$

To finish, we must estimate (6.10). In a similar way with (6.9), we need to estimate

$$\Bigg(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w\rangle_I\Bigg)^{1/2}\Bigg(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(A_I^{w^{-1}})^2|I|\langle w\rangle_I\Bigg)^{1/2}.$$

By Lemma 4.6, we have

$$\begin{aligned}\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(A_I^{w^{-1}})^2|I|\langle w\rangle_I &\le [w^{-1}]_{A_2^d}\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}(A_I^{w^{-1}})^2|I|\langle w^{-1}\rangle_I^{-1}\\ &= [w^{-1}]_{A_2^d}\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\left(\frac{\langle w^{-1}\rangle_{I_+}-\langle w^{-1}\rangle_{I_-}}{\langle w^{-1}\rangle_I^3}\right)^2|I|\\ &\le C[w^{-1}]_{A_2^d}\langle w^{-1}\rangle_J\,.\end{aligned}$$

This completes the proof of Lemma 6.1. $\square$

Due to the almost self adjoint property of the Hilbert transform, a certain bound for π_b^*H immediately returns the same bound for $H\pi_b$. However we have to prove the boundedness of $S\pi_b$ independently because S is not self adjoint.

7 Linear bound for $S\pi_b$

Lemma 7.1. *Let $w\in A_2^d$ and $b\in BMO^d$. Then, there exists C so that*

$$\|S\pi_b\|_{L^2(w)\to L^2(w)}\le C[w]_{A_2^d}\|b\|_{BMO^d}\,.$$

Proof. We are going to prove Lemma 7.1 by showing

$$\langle S\pi_b(w^{-1}f),g\rangle_w\le C[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2(w^{-1})}\|g\|_{L^2(w)}\,, \tag{7.1}$$

for any positive function $f,g\in L^2$. Since $\langle S\pi_b(f),h_I\rangle=\mathrm{sgn}(I)\langle\pi_b(f),h_{\hat I}\rangle=\mathrm{sgn}(I)\langle f\rangle_{\hat I}\langle b,h_{\hat I}\rangle$, We have

$$S\pi_b(f)=\sum_{I\in\mathcal{D}}\mathrm{sgn}(I)\langle f\rangle_{\hat I}\langle b,h_{\hat I}\rangle h_I\,.$$

By expanding g in the disbalanced Haar system for $L^2(w)$,

$$\begin{aligned}\langle S\pi_b(w^{-1}f),g\rangle_w &= \sum_{I\in\mathcal{D}}\langle w^{-1}f\rangle_{\hat I}\langle b,h_{\hat I}\rangle\mathrm{sgn}(I)\langle h_I,g\rangle_w\\ &= \sum_{I\in\mathcal{D}}\sum_{J\in\mathcal{D}}\mathrm{sgn}(I)\langle w^{-1}\rangle_{\hat I}\langle f\rangle_{\hat I,w^{-1}}\langle b,h_{\hat I}\rangle\langle g,h_J^w\rangle_w\langle h_I,h_J^w\rangle_w\,.\end{aligned}$$

Since $\langle h_I,h_J^w\rangle_w$ could be non zero only if $J\supseteq I$, we can split above sum into three parts,

$$\sum_{I\in\mathcal{D}}\mathrm{sgn}(I)\langle w^{-1}\rangle_{\hat I}\langle f\rangle_{\hat I,w^{-1}}\langle b,h_{\hat I}\rangle\langle g,h_I^w\rangle_w\langle h_I,h_I^w\rangle_w, \tag{7.2}$$

$$\sum_{I\in\mathcal{D}}\mathrm{sgn}(I)\langle w^{-1}\rangle_{\hat I}\langle f\rangle_{\hat I,w^{-1}}\langle b,h_{\hat I}\rangle\langle g,h_{\hat I}^w\rangle_w\langle h_I,h_{\hat I}^w\rangle_w, \tag{7.3}$$

and

$$\sum_{I\in\mathcal{D}}\sum_{J:J\supsetneq\hat I}\mathrm{sgn}(I)\langle w^{-1}\rangle_{\hat I}\langle f\rangle_{\hat I,w^{-1}}\langle b,h_{\hat I}\rangle\langle g,h_J^w\rangle_w\langle h_I,h_J^w\rangle_w\,. \tag{7.4}$$

We claim that all sums, (7.2), (7.3), and (7.4), can be bounded with a bound that depends on $[w]_{A_2^d}\|b\|_{BMO^d}$ at most linearly. Since $|\langle h_I, h_I^w\rangle_w| \le \langle w\rangle_I^{1/2}$, we can estimate (7.2)

$$\begin{aligned}
&\Big|\sum_{I\in\mathcal{D}} \operatorname{sgn}(I)\langle w^{-1}\rangle_{\hat I}\langle f\rangle_{\hat I, w^{-1}}\langle b, h_{\hat I}\rangle\langle g, h_I^w\rangle_w\langle h_I, h_I^w\rangle_w\Big|\\
&\qquad\le \Big(\sum_{I\in\mathcal{D}}\langle w^{-1}\rangle_{\hat I}^2\langle f\rangle_{\hat I,w^{-1}}^2\langle b,h_{\hat I}\rangle^2\langle w\rangle_I\Big)^{1/2}\Big(\sum_{I\in\mathcal{D}}\langle g,h_I^w\rangle^2\Big)^{1/2}\\
&\qquad\le C\|g\|_{L^2(w)}[w]_{A_2^d}^{1/2}\Big(\sum_{I\in\mathcal{D}}\langle f\rangle_{I,w^{-1}}^2\langle b,h_I\rangle^2\langle w^{-1}\rangle_I\Big)^{1/2}.
\end{aligned}$$

By Weighted Carleson Embedding Theorem 4.4,

$$\sum_{I\in\mathcal{D}}\langle f\rangle_{I,w^{-1}}^2\langle b,h_I\rangle^2\langle w^{-1}\rangle_I \le C[w]_{A_2^d}\|b\|_{BMO^d}^2\|f\|_{L^2(w^{-1})}^2$$

is provided by

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w^{-1}\rangle_I \le [w]_{A_2^d}\|b\|_{BMO^d}^2\langle w^{-1}\rangle_J$$

which we already have in (6.7). Thus, we have

$$\sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle w^{-1}\rangle_{\hat I}\langle f\rangle_{\hat I,w^{-1}}\langle b,h_{\hat I}\rangle\langle g,h_I^w\rangle_w\langle h_I,h_I^w\rangle_w \le C[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2(w^{-1})}\|g\|_{L^2(w)}. \tag{7.5}$$

Similarly to (7.2), we can estimate (7.3) using $|\langle h_I, h_{\hat I}^w\rangle_w| \le \langle w\rangle_I^{1/2} \le \sqrt{2}\langle w\rangle_{\hat I}^{1/2}$

$$\begin{aligned}
&\Big|\sum_{I\in\mathcal{D}}\operatorname{sgn}(I)\langle w^{-1}\rangle_{\hat I}\langle f\rangle_{\hat I,w^{-1}}\langle b,h_{\hat I}\rangle\langle g,h_{\hat I}^w\rangle_w\langle h_I,h_{\hat I}^w\rangle_w\Big|\\
&\qquad\le \sqrt{2}\sum_{I\in\mathcal{D}}\langle w^{-1}\rangle_{\hat I}\langle f\rangle_{\hat I,w^{-1}}\,|\langle b,h_{\hat I}\rangle|\,\langle g,h_{\hat I}^w\rangle_w\langle w\rangle_{\hat I}^{1/2}\\
&\qquad= 2\sqrt{2}\sum_{I\in\mathcal{D}}\langle w^{-1}\rangle_I\langle f\rangle_{I,w^{-1}}\,|\langle b,h_I\rangle|\,\langle g,h_I^w\rangle_w\langle w\rangle_I^{1/2}\\
&\qquad\le 2\sqrt{2}\Big(\sum_{I\in\mathcal{D}}\langle w^{-1}\rangle_I^2\langle f\rangle_{I,w^{-1}}^2\langle b,h_I\rangle^2\langle w\rangle_I\Big)^{1/2}\Big(\sum_{I\in\mathcal{D}}\langle g,h_I^w\rangle^2\Big)^{1/2}\\
&\qquad\le C\|g\|_{L^2(w)}[w]_{A_2^d}^{1/2}\Big(\sum_{I\in\mathcal{D}}\langle f\rangle_{I,w^{-1}}^2\langle b,h_I\rangle^2\langle w^{-1}\rangle_I\Big)^{1/2}\\
&\qquad\le C[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2(w^{-1})}\|g\|_{L^2(w)}.
\end{aligned}$$

Since h_J^w is constant on $\hat I$, for $J \supsetneq \hat I$ and we denote this constant by $h_J^w(\hat I)$. Then we know by (2.7),

$$\sum_{J: J\supsetneq\hat I}\langle g,h_J^w\rangle_w\langle h_I,h_J^w\rangle_w = \sum_{J:J\supsetneq\hat I}\langle g,h_J^w\rangle_w h_J^w(\hat I)\langle h_I,w\rangle = \langle g\rangle_{\hat I,w}\langle h_I,w\rangle.$$

Thus, we can rewrite (7.4)

$$\begin{aligned}
&\Big|\sum_{I\in\mathcal{D}} \operatorname{sgn}(I)\langle w^{-1}\rangle_{\hat{I}}\langle f\rangle_{\hat{I},w^{-1}}\langle b,h_{\hat{I}}\rangle\langle g\rangle_{\hat{I},w}\langle h_I,w\rangle\Big| \\
&\qquad\le \sum_{I\in\mathcal{D}}\langle w^{-1}\rangle_{\hat{I}}\langle f\rangle_{\hat{I},w^{-1}}\,|\langle b,h_{\hat{I}}\rangle|\,\langle g\rangle_{\hat{I},w}|\langle h_I,w\rangle| \qquad (7.6)\\
&\qquad= \sum_{I\in\mathcal{D}}\langle w^{-1}\rangle_I\,|\langle b,h_I\rangle|\,(|\langle h_{I_-},w\rangle|+|\langle h_{I_+},w\rangle|)\langle f\rangle_{I,w^{-1}}\langle g\rangle_{I,w}\,. \qquad (7.7)
\end{aligned}$$

We claim the sum (7.7) is bounded by $[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2(w^{-1})}\|g\|_{L^2(w)}$. We are going to prove it using Petermichl's Bilinear Embedding Theorem 4.1. Thus, we need to show that the following three embedding conditions hold,

$$\forall\ J\in\mathcal{D},\ \frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle|\,\langle w^{-1}\rangle_I(|\langle h_{I_-},w\rangle|+|\langle h_{I_+},w\rangle|)\frac{1}{\langle w\rangle_I}\le C[w]_{A_2^d}\|b\|_{BMO^d}\,\langle w^{-1}\rangle_J\,, \quad (7.8)$$

$$\forall\ J\in\mathcal{D},\ \frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle|\,\langle w^{-1}\rangle_I(|\langle h_{I_-},w\rangle|+|\langle h_{I_+},w\rangle|)\frac{1}{\langle w^{-1}\rangle_I}\le C[w]_{A_2^d}\|b\|_{BMO^d}\,\langle w\rangle_J\,, \quad (7.9)$$

$$\forall\ J\in\mathcal{D},\ \frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle|\,\langle w^{-1}\rangle_I(|\langle h_{I_-},w\rangle|+|\langle h_{I_+},w\rangle|)\le C[w]_{A_2^d}\|b\|_{BMO^d}\,. \quad (7.10)$$

After we split the sum in (7.8):

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle|\,\langle w^{-1}\rangle_I|\langle h_{I_-},w\rangle|\frac{1}{\langle w\rangle_I}+\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle|\,\langle w^{-1}\rangle_I|\langle h_{I_+},w\rangle|\frac{1}{\langle w\rangle_I}\,,$$

we start with Cauchy-Schwarz inequality to estimate the first sum of embedding condition (7.8),

$$\begin{aligned}
\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle|\,\langle w^{-1}\rangle_I|\langle h_{I_-},w\rangle|\frac{1}{\langle w\rangle_I} &= \frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle|\,\langle w^{-1}\rangle_I\frac{\sqrt{|I_-|}|\Delta_{I_-}w|}{\langle w\rangle_I}\\
&\le\left(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w^{-1}\rangle_I^2\langle w\rangle_I\right)^{1/2}\left(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|I_-||\Delta_{I_-}w|^2\frac{1}{\langle w\rangle_I^3}\right)^{1/2}\\
&\le C[w]_{A_2}^{1/2}\left(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w^{-1}\rangle\right)^{1/2}\left(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|I_-||\Delta_{I_-}w|^2\frac{1}{\langle w\rangle_{I_-}^3}\right)^{1/2}\\
&\le C[w]_{A_2^d}\|b\|_{BMO^d}\langle w^{-1}\rangle_J\,. \qquad (7.11)
\end{aligned}$$

Inequality (7.11) due to Lemma 4.6 and (6.7) . Also, the other sum can be estimated by exactly the same method. Thus we have the embedding condition (7.8). To see the embedding condition (7.9), it is enough to show

$$\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle\,\langle h_{I_-},w\rangle|\le C[w]_{A_2^d}\|b\|_{BMO^d}\langle w\rangle_J\,,$$

as we did above. We use Cauchy-Schwarz inequality for embedding condition (7.9), then

$$
\begin{aligned}
\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle\langle h_{I_-},w\rangle| &= \frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|\langle b,h_I\rangle|\sqrt{|I_-|}|\Delta_{I_-}w| \\
&\leq \left(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\frac{1}{\langle w^{-1}\rangle_I}\right)^{1/2}\left(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|I_-||\Delta_{I_-}w|^2\langle w^{-1}\rangle_I\right)^{1/2} \\
&\leq C\|b\|_{BMO^d}\langle w\rangle_J^{1/2}\left(\frac{[w]_{A_2}}{|J|}\sum_{I\in\mathcal{D}(J)}|I_-||\Delta_{I_-}w|^2\frac{1}{\langle w\rangle_{I_-}}\right)^{1/2} \qquad (7.12) \\
&\leq C[w]_{A_2^d}\|b\|_{BMO^d}\langle w\rangle_J\,. \qquad (7.13)
\end{aligned}
$$

Here inequality (7.12) uses Lemma 4.7, and inequality (7.13) uses the fact that $\langle w\rangle_I^{-1}\leq 2\langle w\rangle_{I_-}^{-1}$ and Theorem 4.5 after shifting the indices.

If we show the embedding condition (7.10), then we can immediately finish the estimate for (7.7) with bound $C[w]_{A_2^d}\|b\|_{BMO^d}\|f\|_{L^2(w^{-1})}\|g\|_{L^2(w)}$. Combining this and (7.5) will give us our desire result. □

8 Proof for embedding condition (7.10)

The following lemma lies at the heart of the matter for the proof of the embedding condition (7.10).

Lemma 8.1. *There is a positive constant C so that for all dyadic interval $J\in\mathcal{D}$*

$$
\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|I|\langle w\rangle_I^{1/4}\langle w^{-1}\rangle_I^{1/4}\left(\frac{|\Delta_{I_+}w|+|\Delta_{I_-}w|}{\langle w\rangle_I}\right)^2\leq C\langle w\rangle_J^{1/4}\langle w^{-1}\rangle_J^{1/4}, \qquad (8.1)
$$

whenever w is a weight. Moreover, if $w\in A_2^d$ then for all $J\in\mathcal{D}$

$$
\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|I|\langle w\rangle_I\langle w^{-1}\rangle_I\left(\frac{|\Delta_{I_+}w|+|\Delta_{I_-}w|}{\langle w\rangle_I}\right)^2\leq C\,[w]_{A_2^d}\,.
$$

Proof of condition (7.10). By using Cauchy-Schwarz inequality and Lemma 8.1, we have:

$$
\begin{aligned}
&\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle\langle w^{-1}\rangle_I(|\langle h_{I_-},w\rangle|+|\langle h_{I_+},w\rangle|) \\
&\quad=\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle\langle w^{-1}\rangle_I\sqrt{\frac{|I|}{2}}\big(|\langle w\rangle_{I_{-+}}-\langle w\rangle_{I_{--}}|+|\langle w\rangle_{I_{++}}-\langle w\rangle_{I_{+-}}|\big) \\
&\quad\leq\frac{1}{\sqrt{2}}\left(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}\langle b,h_I\rangle^2\langle w^{-1}\rangle_I\langle w\rangle_I\right)^{1/2}\left(\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)}|I|\langle w^{-1}\rangle_I\langle w\rangle_I^{-1}(|\Delta_{I_+}w|+|\Delta_{I_-}w|)^2\right)^{1/2} \\
&\quad\leq C[w]_{A_2^d}\|b\|_{BMO^d}\,.
\end{aligned}
$$

□

We turn to the proof of Lemma 8.1. In the first place, we need to revisit some properties of function $B(u,v) := \sqrt[4]{uv}$ on the domain $\mathfrak{D}_0$ which is given by

$$\{(u,v) \in \mathbb{R}^2_+ \,:\, uv \geq 1/2\}\,.$$

It is known, we refer to [2], that $B(u,v)$ satisfies the following differential inequality in $\mathfrak{D}_0$

$$-(du,dv)d^2B(u,v)(du,dv)^t \geq \frac{1}{8}\frac{v^{1/4}}{u^{7/4}}|du|^2\,. \tag{8.2}$$

Furthermore, this implies the following convexity condition. For all (u,v), $(u_\pm, v_\pm) \in \mathfrak{D}_0$,

$$B(u,v) - \frac{B(u_+,v_+) + B(u_-,v_-)}{2} \geq C_1 \frac{v^{1/4}}{u^{7/4}}(u_+ - u_-)^2\,, \tag{8.3}$$

where $u = (u_+ + u_-)/2$ and $v = (v_+ + v_-)/2$. Let us define

$$A(u,v,\Delta u) := aB(u,v) + B(u+\Delta u, v) + B(u - \Delta u, v)\,,$$

on the domain $\mathfrak{D}_1$ with some positive constant $a > 0$. Here $(u,v,\Delta u) \in \mathfrak{D}_1$ means all pairs (u,v), $(u+\Delta u, v)$, $(u-\Delta u, v) \in \mathfrak{D}_0$. Then A has the size property and the convexity property:

$$\text{if } \ (u,v,\Delta u) \in \mathfrak{D}_1, \ \text{ then } \ 0 \leq A(u,v,\Delta u) \leq (a+2)\sqrt[4]{uv}\,, \tag{8.4}$$

and

$$A(u,v,\Delta u) - \frac{1}{2}\big[A(u_+,v_+,\Delta u_1) + A(u_-,v_-,\Delta u_2)\big] \geq C_2 \frac{v^{1/4}}{u^{7/4}}(\Delta u_1^2 + \Delta u_2^2)\,, \tag{8.5}$$

where $u = (u_+ + u_-)/2\,, v = (v_+ + v_-)/2$, and $\Delta u = (u_+ - u_+)/2$. The property (8.5) is directly from the definition of function $B(u,v)$. At the end, Δu will play the role of $\Delta_I w$, Δu_1 is $\Delta_{I_+} w$, and Δu_2 is $\Delta_{I_-} w$. We can rewrite left hand side of the inequality (8.5) as follows

$$\begin{aligned}
&A(u,v,\Delta u) - \frac{1}{2}\big[A(u_+,v_+,\Delta u_1) + A(u_-,v_-,\Delta u_2)\big] \\
&= aB(u,v) + B(u+\Delta u, v) + B(u-\Delta u, v) - \frac{1}{2}\big[aB(u_+,v_+) + B(u_+ + \Delta u_1, v_+) + B(u_+ - \Delta u_1, v_+) \\
&\quad + aB(u_-,v_-) + B(u_- + \Delta u_1, \Delta u_-) + B(u_- - \Delta u_1, v_-)\big] \\
&= aB(u,v) - \frac{a}{2}(B(u_+,v_+) + B(u_-,v_-)) + B(u_+,v) + B(u_-,v) - \frac{1}{2}\Big[B(u+\Delta u + \Delta u_1, v + \Delta v) \\
&\quad + B(u + \Delta u - \Delta u_1, v + \Delta v) + B(u - \Delta u + \Delta u_2, v - \Delta v) + B(u - \Delta u - \Delta u_2, v - \Delta v)\Big]\,.
\end{aligned} \tag{8.6}$$

Using Taylor's theorem:

$$B(u+u_0, v+v_0) = B(u,v) + \nabla B(u,v)(u_0,v_0)^t + \int_0^1 (1-s)(u_0,v_0)d^2B(u+su_0, v+sv_0)(u_0,v_0)^t ds\,,$$

and the convexity condition of $B(u,v)$, we are going to estimate the lower bounds of (8.6).

$$
\begin{aligned}
&-\frac{1}{2}B(u+\Delta u+\Delta u_1, v+\Delta v)\\
&=-\frac{1}{2}\Big(B(u,v)+\nabla B(u,v)(\Delta u+\Delta u_1,\Delta v)^t\Big)\\
&\quad-\int_0^1(1-s)(\Delta u+\Delta u_1,\Delta v)d^2B(u+s(\Delta u+\Delta u_1),v+s\Delta v)(\Delta u+\Delta u_1,\Delta v)^t ds\\
&\ge-\frac{1}{2}\Big(B(u,v)+\nabla B(u,v)(\Delta u+\Delta u_1,\Delta v)^t\Big)+\frac{1}{8}\int_0^1(1-s)\frac{(v+s\Delta v)^{1/4}}{(u+s(\Delta u+\Delta u_1))^{7/4}}(\Delta u+\Delta u_1)^2 ds\\
&\ge-\frac{1}{2}\Big(B(u,v)+\nabla B(u,v)(\Delta u+\Delta u_1,\Delta v)^t\Big)+\frac{(\Delta u+\Delta u_1)^2}{8(4u)^{7/4}}\int_0^1(1-s)(v+s\Delta v)^{1/4}ds \qquad (8.7)\\
&\ge-\frac{1}{2}\Big(B(u,v)+\nabla B(u,v)(\Delta u+\Delta u_1,\Delta v)^t\Big)+\frac{(\Delta u+\Delta u_1)^2v^{1/4}}{8(4u)^{7/4}}\int_0^1(1-s)(1+s\frac{\Delta v}{v})^{1/4}ds\\
&\ge-\frac{1}{2}\Big(B(u,v)+\nabla B(u,v)(\Delta u+\Delta u_1,\Delta v)^t\Big)+\frac{1}{72\cdot 4^{3/4}}\frac{v^{1/4}}{u^{7/4}}(\Delta u+\Delta u_1)^2\,. \qquad (8.8)
\end{aligned}
$$

Inequality (8.7) is due to the following inequalities

$$
|\Delta u|=\frac{|u_+-u_-|}{2}\le\frac{|u_++u_-|}{2}=u \text{ and } |\Delta u_1|=\frac{|u_{++}-u_{+-}|}{2}\le u_+\le 2u\,.
$$

Since $(1-s)^{1/4}\le(1-|\beta|s)^{1/4}\le(1+\beta s)^{1/4}$ for any $|\beta|<1$, it is clear that

$$
\int_0^1(1-s)(1+\beta s)^{1/4}ds\ge\int_0^1(1-s)^{5/4}ds=\frac{4}{9}\,,
$$

and this allows the inequality (8.8). With the same arguments, we also estimate the following lower bounds:

$$
\begin{aligned}
&-\frac{1}{2}\Big[B(u+\Delta u-\Delta u_1,v+\Delta v)+B(u-\Delta u+\Delta u_2,v-\Delta v)+B(u-\Delta u-\Delta u_2,v-\Delta v)\Big]\\
&\quad\ge-\frac{1}{2}\Big(B(u,v)+\nabla B(u,v)(\Delta u-\Delta u_1,\Delta v)^t\Big)+\frac{1}{72\cdot 4^{3/4}}\frac{v^{1/4}}{u^{7/4}}(\Delta u-\Delta u_1)^2\\
&\quad\quad-\frac{1}{2}\Big(B(u,v)+\nabla B(u,v)(-\Delta u+\Delta u_2,-\Delta v)^t\Big)+\frac{1}{72\cdot 4^{3/4}}\frac{v^{1/4}}{u^{7/4}}(-\Delta u+\Delta u_2)^2\\
&\quad\quad-\frac{1}{2}\Big(B(u,v)+\nabla B(u,v)(-\Delta u-\Delta u_2,-\Delta v)^t\Big)+\frac{1}{72\cdot 4^{3/4}}\frac{v^{1/4}}{u^{7/4}}(\Delta u+\Delta u_2)^2\,. \qquad (8.9)
\end{aligned}
$$

We can have the following inequality by combining (8.8), (8.9) and (8.6),

$$
\begin{aligned}
A(u,v,\Delta u)&-\frac{1}{2}\big[A(u_+,v_+,\Delta u_1)+A(u_-,v_-,\Delta u_2)\big]\\
&\ge (a-2)B(u,v)-\frac{a}{2}(B(u_+,v_+)+B(u_-,v_-))+B(u_+,v)+B(u_-,v)\\
&\quad+\frac{1}{36\cdot 4^{3/4}}\frac{v^{1/4}}{u^{7/4}}(2\Delta u^2+\Delta u_1^2+\Delta u_2^2)\\
&\ge \frac{1}{36\cdot 4^{3/4}}\frac{v^{1/4}}{u^{7/4}}(\Delta u_1^2+\Delta u_2^2)\,. \qquad (8.10)
\end{aligned}
$$

To see the inequality (8.10), using convexity condition of $B(u,v)=\sqrt[4]{uv}$ and inequality: $(1-s)u \leq u-s\Delta u \leq u+s\Delta u$,

$$
\begin{aligned}
&(a-2)B(u,v)-\frac{a}{2}\Big(B(u_+,v_+)+B(u_-,v_-)\Big)+B(u_+,v)+B(u_-,v)\\
&\qquad = a\bigg(B(u,v)-\frac{1}{2}(B(u_+,v_+)+B(u_-,v_-)\bigg)\\
&\qquad\quad -\bigg(\frac{3}{16}\Delta u^2\int_0^1(1-s)v^{1/4}(u+s\Delta u)^{-7/4}ds+\frac{3}{16}\Delta u^2\int_0^1(1-s)v^{1/4}(u-s\Delta u)^{-7/4}ds\bigg)\\
&\qquad \geq aC_1\frac{v^{1/4}}{u^{7/4}}\Delta u^2-\frac{6}{16}\Delta u^2\frac{v^{1/4}}{u^{7/4}}\int_0^1(1-s)^{-3/4}ds\\
&\qquad = aC_1\frac{v^{1/4}}{u^{7/4}}\Delta u^2-\frac{3}{2}\Delta u^2\frac{v^{1/4}}{u^{7/4}}\\
&\qquad = \left(aC_1-\frac{3}{2}\right)\frac{v^{1/4}}{u^{7/4}}\Delta u^2\,. \qquad (8.11)
\end{aligned}
$$

Choosing a constant a sufficiently large so that $aC_1>3/2$, quantity in (8.11) remains positive. This observation and discarding nonnegative terms yield inequality (8.10). Choosing the constant $C_2=1/(36\cdot 4^{3/4})$ in (8.5) completes the proof of the concavity property of $A(u,v,\Delta u)$. We now turn to the proof of Lemma 8.1.

Proof of Lemma 8.1. Let $u_I:=\langle w\rangle_I$, $v_I:=\langle w^{-1}\rangle_I$, $u_\pm=u_{I_\pm}$, $v_\pm=v_{I_\pm}$, $\Delta u_I=\Delta_I w$, $\Delta u_1=\Delta u_{I_+}$, and $\Delta u_2=\Delta u_{I_-}$. Then by Hölder's inequality $(u,v,\Delta u)$, $(u_+,v_+,\Delta u_1)$, and $(u_-,v_-,\Delta u_2)$ belong to $\mathfrak{D}_1$. Fix $J\in\mathcal{D}$, by properties (8.4) and (8.5)

$$
\begin{aligned}
(a+2)|J|\sqrt[4]{\langle w\rangle_J\langle w^{-1}\rangle_J} &\geq |J|A(u_J,v_J,\Delta u_J)\\
&\geq \frac{1}{2}\Big(|J_+|A(u_+,v_+,\Delta u_1)+|J_-|A(u_-,v_-,\Delta u_2)\Big)+|J|C\frac{\langle w^{-1}\rangle_J}{\langle w\rangle_J^{7/4}}(|\Delta_{J_+}u|^2+|\Delta_{J_-}u|^2)\,.
\end{aligned}
$$

Since $A(u,v,\Delta u)\geq 0$, iterating the above process will yield

$$|J|\sqrt[4]{\langle w\rangle_J\langle w^{-1}\rangle_J}\geq C\sum_{I\in\mathcal{D}(J)}|I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}(|\Delta_{I_+}w|^2+|\Delta_{I_-}w|^2)\,. \qquad (8.12)$$

Also, one can easily have

$$|J|\sqrt[4]{\langle w\rangle_J\langle w^{-1}\rangle_J}\geq C\sum_{I\in\mathcal{D}(J)}|I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}\,\Delta_{I_+}w^2\,, \qquad (8.13)$$

and

$$|J|\sqrt[4]{\langle w\rangle_J\langle w^{-1}\rangle_J}\geq C\sum_{I\in\mathcal{D}(J)}|I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}\,\Delta_{I_-}w^2\,. \qquad (8.14)$$

Then,

$$
\begin{aligned}
&\frac{1}{|J|}\sum_{I\in\mathcal{D}(J)} |I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}(|\Delta_{I_+}w|+|\Delta_{I_-}w|)^2\\
&\qquad=\frac{1}{|J|}\biggl(\sum_{I\in\mathcal{D}(J)} |I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}(|\Delta_{I_+}w|^2\\
&\qquad\quad+|\Delta_{I_-}w|^2)+2\sum_{I\in\mathcal{D}(J)} |I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}(|\Delta_{I_+}w|\,|\Delta_{I_-}w|)\biggr)\\
&\qquad\le\frac{1}{|J|}\biggl(\sum_{I\in\mathcal{D}(J)} |I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}(|\Delta_{I_+}w|^2+|\Delta_{I_-}w|^2)\\
&\qquad\quad+2\Bigl(\sum_{I\in\mathcal{D}(J)} |I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}\,\Delta_{I_+}w^2\Bigr)^{1/2}\Bigl(\sum_{I\in\mathcal{D}(J)} |I|\langle w^{-1}\rangle_I^{1/4}\langle w\rangle_I^{-7/4}\,\Delta_{I_-}w^2\Bigr)^{1/2}\biggr)\\
&\qquad\le\frac{3}{C}\sqrt[4]{\langle w\rangle_I\langle w^{-1}\rangle_I}\,.
\end{aligned}
$$

□

9 Recently developed tools and their applications

The dyadic shift operator was first introduced in [20] to replace the weighted norm estimate for the Hilbert transform. It was also encountered in [29], so Riesz transforms can be obtained as the result of averaging some dyadic shift operator. Recently, in [17] and [7], a more general class of dyadic shift operators, so called the Haar shift operators were introduced. The Hilbert transform, Riesz transforms, and Beurling-Ahlfors operator are in the convex hull of this class, as they can be written as appropriate averages of Haar shift operators. Let $\mathcal{D}^n$ denote the collection of dyadic cubes in $\mathbb{R}^n$, $\mathcal{D}^n(Q)$ denotes dyadic subcubes of Q, and $|Q|$ denotes the volume of the dyadic cube Q. We start with some definitions.

Definition 9.1. A Haar function on a cube $Q\subset\mathbb{R}^n$ is a function H_Q such that

(a) H_Q is supported on Q, and is constant on $\mathcal{D}^n(Q)$.

(b) $\|H_Q\|_\infty\le|Q|^{-1/2}$.

(c) H_Q has a mean zero.

Definition 9.2. Given an integer $\tau>0$, we say an operator of the following form is in *the first class of Haar shift operators of index* τ

$$
T_\tau f(x)=\sum_{Q\in\mathcal{D}^n}\sum_{\substack{Q',Q''\in\mathcal{D}(Q)\\ 2^{-\tau n}|Q|\le|Q'|,|Q''|}} a_{Q',Q''}\langle f,H_{Q'}\rangle H_{Q''}(x)\,,
$$

where the constant $a_{Q',Q''}$ satisfy the following size condition:

$$
|\,a_{Q',Q''}|\le C\Bigl(\frac{|Q'|}{|Q|}\cdot\frac{|Q''|}{|Q|}\Bigr)^{1/2}. \tag{9.1}
$$

Note that once a choice of Haar functions has been made $\{H_Q\}_{Q\in\mathcal{D}^n}$, then this is an orthogonal family, such that $\|H_Q\|_{L^2} \leq 1$, so one could normalize in L^2 . Note that one can easily see that the dyadic shift operator S belongs to the first class of a Haar shift operator of index $\tau = 1$ with

$$a_{I',I''} = \begin{cases} \pm 1 & \text{for } I' = I, \quad I'' = I_{\mp} \\ 0 & \text{otherwise} \end{cases} .$$

One of the main result in [17] and [7] is the following

Theorem 9.3 ([17], [7]). *Let T be in the first class of Haar shift operators of index τ . Then for all $w \in A_2^d$, there exists $C(\tau, n)$ which only depends on τ and n such that*

$$\|T\|_{L^2(w)\to L^2(w)} \leq C(\tau, n)[w]_{A_2^d} .$$

As a consequence of this Theorem, linear bounds for the Hilbert transform, Riesz transforms, and the Beurling-Ahlfors operator are recovered. There are now two different proofs of Theorem (9.3) in [17] and [7]. The commutator $[\lambda_b, S]$ is also in the first class of Haar shift operators of index $\tau = 1$. Recall the observation in the section 4,

$$[\lambda_b, S](f) = -\sum_{I\in\mathcal{D}} \Delta_I b \langle f, h_I\rangle (h_{I_+} + h_{I_-}) .$$

Then we can see

$$a_{I',I''} = \begin{cases} -\Delta_I b & \text{for } I' = I, \quad I'' = I_{\pm} \\ 0 & \text{otherwise} \end{cases} ,$$

moreover $|\, a_{I',I''}| = |\Delta_I b| \leq 2\|b\|_{BMO^d}$ this means the constant $a_{I',I''}$ satisfy the size condition (9.1) with $C = 2\sqrt{2}\|b\|_{BMO^d}$. These observations, Theorem 5.1, and Theorem 9.3 immediately recover the quadratic bound for the commutator of the Hilbert transform which was proved in this paper. We now define the second class of Haar shift operators of index τ .

Definition 9.4. Given an integer $\tau > 0$, we say an operator T of the form in Definition 9.2 is in *the second class of Haar shift operators of index τ* , if T is bounded on L^2 and the function H_Q satisfy the condition (a) and (b) in Definition 9.2.

The second class of Haar shift operators is more general than the first class. One can easily observe that the operators π_b, $S\pi_b$ and $\pi_b^* S$ do not satisfy the condition (c) on Definition 9.1, however these operators satisfy the conditions of Definition 9.4. Note that the n-variable paraproduct is a sum of $2^n - 1$ of the second class of Haar shift operator of index 1, the restricted n-variable dyadic paraproduct

$$\pi_b f = \sum_{Q\in\mathcal{D}^n} \langle f\rangle_Q \langle b, H_Q\rangle H_Q .$$

In [14], the linear estimate for the maximal truncations of these operators is presented. This also recovers our linear bound estimates for $S\pi_b$ and $\pi_b^* S$. On the other hand, authors in [7] also reproduce the linear estimate for the dyadic paraproduct with different technique.

Lemma 9.5. *Let T_τ a Haar shift operator of the first class, then $[\lambda_b, T_\tau]$ is an operator of the same class.*

Proof. We are going to use the restricted multi-variable λ_b operator which is

$$\lambda_b f = \sum_{Q\in\mathcal{D}^n} \langle b\rangle_Q \langle f, H_Q\rangle H_Q .$$

One can get the n-variable λ_b operator by summing over $2^n - 1$ of restricted λ_b operator. Observe that,

$$\begin{aligned}
[\lambda_b, T_\tau]f &= \lambda_b(T_\tau f) - T_\tau(\lambda_b f) \\
&= \sum_{Q\in\mathcal{D}_n} \sum_{\substack{Q',Q''\in\mathcal{D}(Q) \\ 2^{-\tau n}|Q|\le|Q'|,|Q''|}} a_{Q',Q''}\langle b\rangle_{Q''}\langle f, H_{Q'}\rangle H_{Q''} - \sum_{Q\in\mathcal{D}_n} \sum_{\substack{Q',Q''\in\mathcal{D}(Q) \\ 2^{-\tau n}|Q|\le|Q'|,|Q''|}} a_{Q',Q''}\langle b\rangle_{Q'}\langle f, H_{Q'}\rangle H_{Q''} \\
&= \sum_{Q\in\mathcal{D}_n} \sum_{\substack{Q',Q''\in\mathcal{D}(Q) \\ 2^{-\tau n}|Q|\le|Q'|,|Q''|}} a_{Q',Q''}(\langle b\rangle_{Q''} - \langle b\rangle_{Q'})\langle f, H_{Q'}\rangle H_{Q''}\,.
\end{aligned}$$

Since

$$\left|\, a_{Q',Q''}(\langle b\rangle_{Q''} - \langle b\rangle_{Q'})\right| \le \|b\|_{BMO}\left|\, a_{Q',Q''}\right|,$$

$[\lambda_b, T_\tau]$ remains in the same class of T_τ . □

Theorem 9.3 and Lemma 9.5 allow to extend our result to more general class of commutators including the Riesz transforms and the Beurling-Ahlfors operator as in Theorem 1.3.

10 Sharp bounds

In this section, we start proving that the quadratic estimate in Theorem 1.5 is sharp, by showing an example which returns quadratic bound. This example was discovered by C. Peréz [22] who is kindly allowing us to reproduce it in this paper. The same calculations show that the bounds in Theorem 1.1 are also sharp for $p \neq 2$ and $1 < p < \infty$. Variations over this example will then show that the bounds in Theorem 1.3 are sharp for the Riesz transforms and the Beurling-Ahlfors operator as well.

10.1 The Hilbert transform

Consider the weight, for $0 < \delta < 1$:

$$w(x) = |x|^{1-\delta}\,.$$

It is well known that w is an A_2 weight and

$$[w]_{A^2} \sim \frac{1}{\delta}\,.$$

We now consider the function $f(x) = x^{-1+\delta}\chi_{(0,1)}(x)$ and BMO function $b(x) = \log|x|$. We claim that

$$|[b,H]f(x)| \ge \frac{1}{\delta^2} f(x)\,.$$

For $0 < x < 1$, we have

$$\begin{aligned}
[b,H]f(x) &= \int_0^1 \frac{\log x - \log y}{x-y} y^{-1+\delta} dy = \int_0^1 \frac{\log(x/y)}{x-y} y^{-1+\delta} dy \\
&= x^{-1+\delta} \int_0^{1/x} \frac{\log\left(\frac{1}{t}\right)}{1-t} t^{-1+\delta} dt\,.
\end{aligned}$$

Now,

$$\int_0^{1/x} \frac{\log(1/t)}{1-t} t^{-1+\delta} dt = \int_0^1 \frac{\log(1/t)}{1-t} t^{-1+\delta} dt + \int_1^{1/x} \frac{\log(1/t)}{1-t} t^{-1+\delta} dt \,,$$

and since $\frac{\log(1/t)}{1-t}$ is positive for $(0,1) \cup (1,\infty)$ we have for $0 < x < 1$

$$|[b,H]f(x)| > x^{-1+\delta} \int_0^1 \frac{\log(1/t)}{1-t} t^{-1+\delta} dt \,. \tag{10.1}$$

But since

$$\int_0^1 \frac{\log(1/t)}{1-t} t^{-1+\delta} dt > \int_0^1 \log(1/t) t^{-1+\delta} dt = \int_0^\infty s e^{-s\delta} ds = \frac{1}{\delta^2} \,, \tag{10.2}$$

our claim follows and

$$\|[b,H]f\|_{L^2(w)} \geq \frac{1}{\delta^2} \|f\|_{L^2(w)} \sim [w]_{A_2}^2 \|f\|_{L^2(w)} \,.$$

A first approximation of what the bounds in $L^p(w)$ is given by an application of the sharp extrapolation theorem for the upper bound, paired with the knowledge of the sharp bound on $L^2(w)$ to obtain a lower bound.

Proposition 10.1. *For $1 < p < \infty$ there exist constants c and C only depending on p such that*

$$c[w]_{A_p}^{2\min\{1,\frac{1}{p-1}\}} \|b\|_{BMO} \leq \|[b,H]\|_{L^p(w)\to L^p(w)} \leq C[w]_{A_p}^{2\max\{1,\frac{1}{p-1}\}} \|b\|_{BMO} \,, \tag{10.3}$$

for all $b \in BMO$.

Proof. Because the upper bound in (10.3) is the direct consequence of the quadratic bound in the Theorem 1.5 and sharp extrapolation theorem, we will only prove the lower bound. Let us assume that, for $1 < r < 2$ and $\alpha < 1$,

$$\|[b,H]\|_{L^r(w)\to L^r(w)} \leq C[w]_{A_r}^{2\alpha} \|b\|_{BMO} \,.$$

This and the sharp extrapolation theorem return

$$\|[b,H]\|_{L^2(w)\to L^2(w)} \leq C[w]_{A_2}^{2\alpha} \|b\|_{BMO} \,.$$

This contradicts to the sharpness $(p = 2)$. Similarly, one can conclude for $p > 2$. ☐

We now consider the weight $w(x) = |x|^{(1-\delta)(p-1)}$ then w is an A_p weight with $[w]_{A_p} \sim \delta^{1-p}$. By (10.1) and (10.2) we have

$$\|[b,H]f\|_{L^p(w)} \geq \frac{1}{\delta^2} \|f\|_{L^p(w)} = (\delta^{1-p})^{\frac{2}{p-1}} \|f\|_{L^p(w)} \sim [w]_{A_p}^{\frac{2}{p-1}} \|f\|_{L^p(w)} \,.$$

This shows the upper bound in (10.3) is sharp for $1 < p \leq 2$. We use the duality argument to see the sharpness of the quadratic estimate for $p > 2$. Note that the commutator is a self-adjoint operator:

$$\langle bH(f) - H(bf), g\rangle = \langle f, H^*(bg)\rangle - \langle f, bH^*(g)\rangle = \langle f, bH(g) - H(bg)\rangle \,.$$

Consider $1 < p \leq 2$ and set $u = w^{1-p'}$, then

$$\begin{aligned} \|[b,H]\|_{L^{p'}(u)\to L^{p'}(u)} &= \|[b,H]\|_{L^{p'}(w^{1-p'})\to L^{p'}(w^{1-p'})} = \|[b,H]^*\|_{L^{p'}(w^{1-p'})\to L^{p'}(w^{1-p'})} \\ &= \|[b,H]\|_{L^p(w)\to L^p(w)} \leq C\|b\|_{BMO^d}[w]_{A_p}^{\frac{2}{1-p}} \\ &= C\|b\|_{BMO}[w^{1-p'}]_{A_{p'}}^2 = C\|b\|_{BMO}[u]_{A_{p'}}^2 \,. \end{aligned} \tag{10.4}$$

Since the inequality in (10.4) is sharp, we can conclude that the result of Theorem 1.1 is also sharp for $p > 2$.

10.2 Beurling-Ahlfors operator

Recall the Beurling-Ahlfors operator $\mathcal{B}$ is given by convolution with the distributional kernel $p.v.1/z^2$:

$$\mathcal{B}f(x,y) = p.v.\frac{1}{\pi}\int_{\mathbb{R}^2}\frac{f(x-u,y-v)}{(u+iv)^2}\,dudv\,.$$

Then the commutator of the Beurling-Ahlfors operator can be written:

$$[b,\mathcal{B}]f(x,y) = p.v.\frac{1}{\pi}\int_{\mathbb{R}^2}\frac{b(x,y)-b(s,t)}{((x-s)+i(y-t))^2}\,f(s,t)\,dsdt\,.$$

It was observed, in [9], that the linear bound for the Beurling-Ahlfors operator is sharp in $L^2(w)$, with weights $w(z) = |\,z|^{\alpha}$ and functions $f(z) = |\,z|^{-\alpha}$ where $|\,\alpha| < 2$. Similarly, we consider weights $w(z) = |\,z|^{2-\delta}$ where $0<\delta<1$. Note that $w(z) = |\,z|^{2-\delta} : \mathbb{C}\to[0,\infty)$ is a A_2-weight with $[w]_{A_2}\sim\delta^{-1}$. We also consider a BMO function $b(x) = \log|\,z|$. Let $E = \{(r,\theta)\,|\,0<r<1,\,0<\theta<\pi/2\}$ and $\Omega = \{(r,\theta)\,|\,1<r<\infty,\,\pi<\theta<3\pi/2\}$ We are going to estimate $|[b,\mathcal{B}]f(z)|$ for $z\in\Omega$ with a function $f(z) = |\,z|^{\delta-2}\chi_E(z)$. Let $z = x+iy$ and $\zeta = s+ti$. Then, for $z\in\Omega$,

$$\begin{aligned}
|[b,\mathcal{B}]f(z)| &= \frac{1}{\pi}\left|\int_E\frac{(b(z)-b(\zeta))f(\zeta)}{(z-\zeta)^2}\,d\zeta\right| \\
&= \frac{1}{\pi}\left|\int_E\frac{b(x,y)-b(s,t)}{((x-s)+i(y-t))^2}f(s,t)dsdt\right| \\
&= \frac{1}{\pi}\left|\int_E\frac{(\log|\,z|-\log|\,\zeta|)|\,\zeta|^{\delta-2}((x-s)^2-(y-t)^2)}{((x-s)^2+(y-t)^2)^2}\,dsdt\right. \\
&\qquad\left.+\,i\int_E\frac{(\log|\,z|-\log|\,\zeta|)|\,\zeta|^{\delta-2}(2(x-s)(y-t))}{((x-s)^2+(y-t)^2)^2}\,dsdt\right|.
\end{aligned}$$

For $z\in\Omega$ and $\zeta\in E$, we have $(x-s)(y-t)\geq xy$ and by triangle inequality $((x-s)^2+(y-t)^2)^2 = |\,z-\zeta|^4\leq(|\,z|+|\,\zeta|)^4$. After neglecting the positive term (real part), we get

$$\begin{aligned}
|[b,\mathcal{B}]f(z)|^2 &\geq \frac{4}{\pi^2}\left(xy\int_E\frac{\log(|\,z|/|\,\zeta|)|\,\zeta|^{\delta-2}}{(|\,z|+|\,\zeta|)^4}\,dsdt\right)^2 = \frac{4}{\pi^2}\left(xy\int_0^{\pi/2}\int_0^1\frac{\log(|\,z|/r)r^{\delta-2}r}{(|\,z|+r)^4}\,drd\theta\right)^2 \\
&= x^2y^2\left(\frac{1}{|\,z|^4}\int_0^1\frac{\log(|\,z|/r)r^{\delta-1}}{(1+r/|\,z|)^4}dr\right)^2 = x^2y^2\left(\frac{1}{|\,z|^4}\int_0^{1/|z|}\frac{\log(1/t)(|\,z|t)^{\delta-1}}{(1+t)^4}\,|\,z|dt\right)^2 \\
&= x^2y^2\left(\frac{1}{|\,z|^{4-\delta}}\int_0^{1/|z|}\frac{\log(1/t)t^{\delta-1}}{(1+t)^4}\,dt\right)^2.
\end{aligned}$$

Since $|\,z|/(|\,z|+1)\leq 1/(1+t)$, for $t<1/|\,z|$, we have

$$\begin{aligned}
|[b,\mathcal{B}]f(z)|^2 &\geq x^2y^2\left(\frac{1}{|\,z|^{-\delta}(|\,z|+1)^4}\int_0^{1/|z|}\log(1/t)t^{\delta-1}\,dt\right)^2 \\
&= \frac{x^2y^2}{|\,z|^{-2\delta}(|\,z|+1)^8}\left(\frac{|\,z|^{-\delta}(1+\delta\log|\,z|)}{\delta^2}\right)^2 \\
&= \frac{x^2y^2}{(|\,z|+1)^8}\frac{(1+\delta\log|\,z|)^2}{\delta^4}\,.
\end{aligned}$$

Then, we can estimate the $L^2(w)$-norm as follows.

$$
\begin{aligned}
\|[b,\mathcal{B}]f\|^2_{L^2(w)} &\geq \frac{1}{\delta^4}\int_\Omega \frac{x^2y^2(1+\delta\log|z|)^2}{(|z|+1)^8}\,|z|^{2-\delta}\,dxdy \\
&= \frac{1}{\delta^4}\int_1^\infty\int_\pi^{3\pi/2}\frac{r^4\cos^2\theta\sin^2\theta(1+\delta\log r)^2}{(r+1)^8}\,r^{3-\delta}drd\theta \\
&= \frac{\pi}{\delta^4 16}\int_1^\infty \frac{r^{7-\delta}(1+\delta\log r)^2}{(r+1)^8}\,dr \geq \frac{\pi}{\delta^4 16}\int_1^\infty \frac{r^{7-\delta}(1+\delta\log r)^2}{(2r)^8}\,dr \\
&= \frac{\pi}{\delta^4 2^{12}}\int_1^\infty (1+\delta\log r)^2 r^{-1-\delta}\,dr \\
&= \frac{\pi}{\delta^4 2^{12}}\left(\frac{1}{\delta}+\frac{2\delta}{\delta^2}+\frac{2\delta^2}{\delta^3}\right) = \frac{5\pi}{2^{12}}\cdot\frac{1}{\delta^5}\,.
\end{aligned}
$$

Combining with $\|f\|^2_{L^2(w)} = \pi/2\delta$, we have that $\|[b,\mathcal{B}]f\|_{L^2(w)}/\|f\|_{L^2(w)} \sim \delta^{-2}$, which allows to conclude that the quadratic bound for the commutator with the Beurling-Ahlfors operators is sharp in $L^2(w)$. Same calculations with weights $w(z) = |z|^{(2-\delta)(p-1)}$ and functions $f(z) = |z|^{(\delta-2)(p-1)}$ will provide the sharpness for $1 < p \leq 2$, and it is sufficient to conclude for all $1 < p < \infty$ because the Beurling-Ahlfors operator is essentially self adjoint operator ($\mathcal{B}^* = e^{i\phi}\mathcal{B}$), so the commutator of the Beurling-Ahlfors operator is also self adjoint.

10.3 Riesz transforms

Consider weights $w(x) = |x|^{n-\delta}$ and functions $f(x) = x^{\delta-n}\chi_E(x)$ where $E = \{x\,|\,x\in(0,1)^n \cap B(0,1)\}$, and a BMO function $b(x) = \log|x|$. It was observed that $|x|^{n-\delta}$ is an A_2-weight in $\mathbb{R}^n$ with $[w]_{A^2} \sim \delta^{-1}$. We are going to estimate $[b,R_j]f$ over the set $\Omega = \{y \in B(0,1)^c\,|\,y_i < 0 \text{ for all } i = 1,2,\ldots,n\}$, where R_j stands for the j-th direction Riesz transform on $\mathbb{R}^n$ and is defined as follows:

$$
R_jf(x) = c_n p.v.\int_{\mathbb{R}^n}\frac{y_j}{|y|^{n+1}}\,f(x-y)\,dy\,, \qquad 1 \leq j \leq n\,,
$$

where $c_n = \Gamma((n+1)/2)/\pi^{(n+1)/2}$. One can observe that, for all $x \in E$ and fixed $y \in \Omega$,

$$
|y_j - x_j| \geq |y_j| \text{ and } |y-x| \leq |y| + |x|\,.
$$

C_n denotes a constant depending only on the dimension that may change from line to line. Then,

$$
\begin{aligned}
|[b,R_j]f(y)| &= \left|\int_E \frac{(y_j-x_j)(\log|y|-\log|x|)|x|^{\delta-n}}{|y-x|^{n+1}}\,dx\right| \geq |y_j|\int_E \frac{\log(|y|/|x|)|x|^{\delta-n}}{(|y|+|x|)^{n+1}}\,dx \\
&\geq |y_j|\int_{E\cap S^{n-1}}\int_0^1 \frac{\log(|y|/r)r^{\delta-n}r^{n-1}}{(|y|+r)^{n+1}}\,drd\sigma = C_n|y_j|\int_0^{1/|y|}\frac{\log(1/t)(t|y|)^{\delta-1}|y|}{(|y|+|y|t)^{n+1}}\,dt \\
&= \frac{C_n|y_j|}{|y|^{n+1-\delta}}\int_0^{1/|y|}\frac{\log(1/t)t^{\delta-1}}{(1+t)^{n+1}}\,dt \geq \frac{C_n|y_j|}{|y|^{n+1-\delta}}\left(\frac{|y|}{|y|+1}\right)^{n+1}\int_0^{1/|y|}\log(1/t)t^{\delta-1}\,dt \\
&= \frac{C_n|y_j|}{|y|^{-\delta}(|y|+1)^{n+1}}\left(\frac{|y|^{-\delta}(1+\delta\log|y|)}{\delta^2}\right) = \frac{C_n|y_j|}{(|y|+1)^{n+1}}\frac{1+\delta\log|y|}{\delta^2}\,.
\end{aligned}
$$

We now can bound from below the $L^2(w)$-norm as follows.

$$
\begin{aligned}
\|[b,R_j]f(x)\|^2_{L^2(w)} &> \frac{C_n}{\delta^4}\int_{\Omega}\frac{y_j^2(1+\delta\log|\,y|)^2}{(|\,y|+1)^{2n+2}}\,|\,y|^{n-\delta}dy \\
&\geq \frac{C_n}{\delta^4}\int_{\Omega\cap S^{n-1}}\int_1^{\infty}\frac{\gamma_j^2r^2(1+\delta\log r)^2r^{n-\delta}r^{n-1}}{(r+1)^{2n+2}}\,drd\sigma(\gamma) \\
&\geq \frac{C_n}{\delta^4}\int_1^{\infty}\frac{(1+\delta\log r)^2r^{2n-\delta+1}}{r^{2n+2}}\,dr \\
&= \frac{C_n}{\delta^4}\int_1^{\infty}(1+\delta\log r)^2r^{-\delta-1}\,dr = \frac{C_n}{\delta^5}\,,
\end{aligned}
$$

which establishes sharpness for the commutator of the Riesz transform when $p = 2$. Since $R_j^* = -R_j$, one can easily check that the commutator of Riesz transforms are also self-adjoint operators. Furthermore, choosing weight $w(x) = |\,x|^{(n-\delta)(p-1)}$, we will obtain the sharpness for $1 < p < \infty$ by the same argument we used in the case of the Hilbert transform.

11 Acknowledgments

This work is part of the author's Ph.D dissertation [4]. The author like to thank Professor Carlos Pérez for his comments and useful suggestions. Finally, the author is also very grateful to his graduate adviser María Cristina Pereyra for her suggestions and helpful interaction.

References

[1] J. Alvarez, R. J. Bagby, D. S. Kurtz and C. Pérez, *Weighted estimates for commutators of linear operators* Studia Math. 104(2) (1993) 195-209.

[2] O. Beznosova, *Linear bound for dyadic paraproduct on weighted Lebesgue space* $L^2(w)$ J. of Fun. Anal. 255 4 (2008), 994-1007.

[3] S. Buckley, *Summation condition on Weights,* Michigan Math. J., 40(1) (1993) 153-170.

[4] D. Chung, *Ph.D Dissertation,* University of New Mexico, (2010).

[5] R. Coifman and C. Fefferman, *Weighted norm inequalities for maximal functions and singular integrals* Studia Math. 51 (1974), 241-250.

[6] R. Coifman, R. Rochberg, and G. Weiss, *Factorization theorems for Hardy spaces in several variables* Ann. of Math. (2) 103 (1976), 611-635.

[7] D. Cruz-Uribe, SFO, J. Martell and C. Pérez, *Sharp weighted estimates for classical operators* Preprint.

[8] O. Dragicević, L. Grafakos, M. C. Pereyra, and S. Petermichl, *Extrapolation and sharp norm estimates for classcical operators on weighted Lebesgue spaces* Publ. Mat. 49 (2005), 73-91.

[9] O. Dragicević and A. Volberg, *Sharp estimate of the Ahlfor-Beurling operator via averaging Martingale transforms*, Michigan Math. J. 51 (2003)

[10] J. B. Garnett, *Bounded analytic functions* Acad. Press, NY, 1981.

[11] H. Helson and G. Szegö, *A problem in prediction theory,* Ann. Math. Pura. Appl. 51 (1960), 107-138.

[12] S. Hukovic, S. Treil, A. Volberg, *The Bellman functions and the sharp weighted inequalities for square functions* Operator Theory : Advances and Applications, the volume in memory of S. A. Vinogradov, v. 113, Birkhauser Verlag, (2000)

[13] R. Hunt, B. Muckenhoupt, and R. Wheeden, *Weighted norm inequalities for conjugate function and the Hilbert transform* Trans. Amer. Math. Soc. 176 (1973), 227-251.

[14] T. Hytönen, M. Lacey, M. Reguera, and A. Vagharshakyan, *Weak and Strong-type estimates for Haar Shift Operators: Sharp power on the A_p characteristic* (2009), available at http://arxiv.org/abs/0911.0713

[15] M. Lacey, *Haar shifts, Commutators, and Hankel operators* CIMPA-UNESCO conference on Real Analysis and it Applications, La Falda Argentina, (2008).

[16] M. Lacey, S. Petermichl, J. Piper and B. Wick, *Iterated Riesz Commutators: A simple proof of boundedness* Proc. of El Escorial (2008).

[17] M. Lacey, S. Petermichl and M. Reguera, *Sharp A_2 inequality for Haar shift operators* Math. Ann. to appear.

[18] B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function* Trans. Amer. Math. Soc. 165 (1972), 207-226.

[19] F. Nazarov, S. Treil, and A. Volberg, *The Bellman functions and two-weight inequalities for Haar multipliers* J. Amer. Math. Soc. 12 (1999), 909-928.

[20] C. Pérez, *Endpoint estimates for commutators of singular integral operators* J. Funct. Anal. 128 (1995), 163-185.

[21] C. Pérez, *Sharp estimats for commutators of singular integrals via iterations of the Hardy-Littlewood maximal functions* J. Fourier Anal. Appl. 3 (1997), 743-756.

[22] C. Pérez, *Personal communication* (2009).

[23] C. Pérez and G. Pradolini, *Sharp weighted Endpoint Estimates for commutators of singular integrals* Michigan Math. H. 49 (2001) 23-37.

[24] M. C. Pereyra, *Haar Multipliers meet Bellman Functions* Revista Mat. Iber. 25 3 (2009), 799-840.

[25] S. Petermichl, *Dyadic shifts and a logarithmic estimate for Hankel operators with matrix symbol* C. R. Acad. Sci. Paris 330 (2000), 455-460.

[26] S. Petermichl, *The sharp bound for the Hilbert transform on weighted Lebesgue spaces in terms of the classical A_p Characteristic,* Amer. J. of Math. 129 (2007) 1355-1375.

[27] S. Petermichl, *The sharp weighted bound for the Riesz transforms* Proc. Amer. Math. Soc. 136 (2008), 1237-1249.

[28] S. Petermichl and S. Pott, *An estimate for weighted Hilbert transform via square functions* Trans. Amer. Math. Soc. 354 (2002), 1699-1703.

[29] S. Petermichl, S. Treil and A. Volberg, *Why the Riesz transforms are averages of the dyadic shifts?* Proc. of the 6th Inter. Conf. on Harmonic Analysis and Partial Differential Equations (El Escorial, 2000) (2002) 209-228.

[30] S. Petermichl and A. Volverg, *Heating of the Ahlfors-Beurling operator: Weakly quasiregular maps on the plane are quasiregular* Duke Math J. 112 (2002), 281-305.

[31] A. Vagharshakyan *Recovering Singular Integrals from Haar Shifts* (2009), available at http://arxiv.org/abs/0911.4968

[32] J. Wittwer, *A shap estimate on the norm of Martingale transform* Math. Res. Lett. 7 (2000) 1-12.

DEPARTMENT OF MATHEMATICS AND STATISTICS, 1 UNIVERSITY OF NEW MEXICO, ALBUQUERQUE, NM 87131-0001, *E-mail:* dwchung@unm.edu and chdaewon@gmail.com